

8378-53

04-09-2010

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Robert F. Bischoff
Agent for Knox Fertilizer Company, Inc
C/O Total Turf Consulting LLC
26 A. A Cooper Rd
Meigs, GA 31765

APR 9 2010

Subject: Label Notification(s) for Pesticide Registration Notice 2007-1

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-1 dated March 23, 2010 for:

EPA Registration 8378-53 Dimension 130 Turf Herbicide with Plant Food

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-1 and finds that the label change(s) requested falls within the scope of PRN-2007-1. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 8378- 53	2. EPA Product Manager Kathryn Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dimension 130 Turf Herbicide with Plant Food	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Knox Fertilizer Company, Inc. P. O. Box 248 Knox, IN 46534 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. <u>N/A</u> Product Name <u>N/A</u>	

Section - II

Amendment - Explain below. Final printed labels in response to Agency letter dated _____

Resubmission in response to Agency letter dated _____ "Me Too" Application.

Notification - Explain below. Other - Explain below.

NOTIFICATION
APR 9 - 2010

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change relative to PR Notice 2007-1. This notification is consistent with the guidance in PR Notice 2007-1 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec.1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 2007-1 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input checked="" type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container
15; 18; 20; 36; 40; 50; 54; 60 lbs

5. Location of Label Directions
 on label

6. Manner in Which Label is Affixed to Product
 Lithograph Paper glued Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Robert F. Bischoff	Title Agent, TotalTurf Consulting LLC	Telephone No. (Include Area Code) (317) 873-0222
----------------------------	--	---

Certification
 I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature <i>Robert F. Bischoff</i>	3. Title Agent, TotalTurf Consulting LLC	6. Date Application Received (Stamped)
4. Typed Name Robert F. Bischoff	5. Date March 23, 2010	

3
11

**TotalTurf Consulting LLC
26 A. A. Cooper Rd.
Meigs, GA 31765**

**Direct: 843-384-2807
Fax: 229-683-3426**

E-Mail: rbcooper@juno.com

March 23, 2010

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7505P)
U. S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attn: Linda Arrington

Subject: Dimension 130 Turf Herbicide with Plant Food
EPA Reg. No. 8378-53
Notification per PR Notice 2007-1
Removal of Toll Free Number from the Disposal Instructions Section


As allowed by PR Notice 2007-1 (Section II.C.3.), the subject product label has been revised to show the removal of the statement "or 1-800-CLEANUP" from the Disposal Instructions Section. The revised label is being submitted by notification on behalf of Knox Fertilizer Company, P. O. Box 248, Knox, IN 46534 by TotalTurf Consulting LLC, an authorized agent for Knox Fertilizer Company.

The following documents are included in the subject submission:

- Cover Letter
- Application for Pesticide (EPA Form No. 8570-1)
- One Copy of Highlighted Final Printed Label (Coded as 20100323.Final Label (PRN2007-1))
(Commercial)
- One Copy of Highlighted Final Printed Label (Coded as 20100323.Final Label (PRN2007-1)) (Retail)

If there are any questions or additional information is needed concerning this submission, do not hesitate to contact me at (317) 873-0222 or Raymond B. Cooper at (843) 384-2807.

Sincerely,



Robert F. Bischoff
Agent
TotalTurf Consulting LLC
26 A. A. Cooper Road
Meigs, GA 31765

4
11

Dimension® 130

Turf Herbicide with Plant Food

This product is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

NOTIFICATION

APR 9 - 2010

In New York State:
For Use Only by Professional Turf Applicators
Use is Prohibited in Nassau and Suffolk Counties

ACTIVE INGREDIENT:

Dithiopyr* 0.13%

INERT INGREDIENTS: **99.87%**

Total 100.00%

*CAS # 97886-45-8

Fertilizer Guarantee Analysis

Nitrogen XX%
Phosphate XX%
Potash XX%
Derived from Urea, Ammoniated Phosphate, and Muriate of Potash.

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA REG. NO. 8378-53

EPA EST. NO. 8378-IN-1

Read the entire label before using this product.
Use only according to label instructions.

NOTICE: Before using this product, read the entire Precautionary Statements, Terms and Conditions of Use, Directions for Use, Use Precautions, and Storage and Disposal Instructions. If the Terms and Conditions of Use are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

Net Weight XX Pounds (XX kg)

Dimension is a registered trademark of Dow AgroSciences LLC

-----End Front Panel-----

5
11

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 from 6:30 a.m. to 4:30 p.m. Pacific Time, everyday, for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS and DOMESTIC ANIMALS
CAUTION: CAUSES EYE IRRITATION. HARMFUL IF INHALED.
 Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

This product should only be applied to lawns and ornamental turfs that are composed of one or more of the following turfgrass species that have been determined to be tolerant. When applied as directed by this label the following established turfgrasses are tolerant to this product:

Cool-Season Grasses	
Common Name	Scientific Name
Bentgrass, Creeping*	Agrostis palustris
Bluegrass, Kentucky	Poa pratensis
Fescue, fine**	Festuca rubra
Fescue, tall	Festuca arundinacea
Ryegrass, perennial	Lolium perenne

Warm-Season Grasses	
Common Name	Scientific Name
Bahiagrass	Paspalum notatum
Bermudagrass	Cynodon dactylon
Buffalograss***	Buchloe dactyloides
Carpetgrass	Axonopus affinis
Centipedegrass	Eremochloa ophiuroides
Kikuyugrass	Pennisetum clandestinum
St. Augustinegrass	Stenotaphrum secundatum
Zoysiagrass	Zoysia japonica

DO NOT apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties. *Use of this product on certain varieties of creeping bentgrass such as 'Cohansey', 'Carmer', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. ** Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested. *** DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the phytotoxicity of this product to other turfgrass varieties on a limited area, at recommended use rates, prior to initiating large-scale use.

Reseeding, Overseeding, or Sprigging
 Reseeding, overseeding, or sprigging of treated areas within 12 weeks after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization must be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

Use Precautions
Turfgrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

- DO NOT use this product on golf course putting greens.
- The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.
- For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.
- Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.
- DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.
- DO NOT graze treated areas.
- Keep people and pets off treated areas until dust has settled.

Control of Crabgrass, Preemergence and Early Postemergence Control
This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make an application and still control crabgrass.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 8.83 lbs. of this product per 1,000 sq. ft. per application, and no more than 13.4 lbs. of this product per 1,000 sq. ft. per year.

In the State of New York do not exceed 0.5 lbs. active ingredient per acre per year (8.8 lbs./1,000 sq. ft.). Use is prohibited in Nassau and Suffolk Counties.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the two control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program #1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program #2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at site where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.


Where split fertilizer applications are recommended the rates in Table 1 may be split between two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1: Recommended Single Application Use Rates*
(Pounds of this product/1,000 sq. ft.)

PROGRAM	#1	#2
USE (turfgrass cut)	Preemergence (high-cut turf)	Preemergence (low-cut turf) Postemergence (low-cut turf)
REGION		
North	2.20 lbs./1,000 sq. ft.	4.41 lbs./1,000 sq. ft.
Transition	3.17 lbs./1,000 sq. ft.	6.53 lbs./1,000 sq. ft.
South	3.37 lbs./1,000 sq. ft.	8.83 lbs./1,000 sq. ft.
Coastal South	6.53 lbs./1,000 sq. ft.	8.83 lbs./1,000 sq. ft.**
*DO NOT apply more than 8.83 lbs. of this product/1,000 sq. ft. per application, and no more than 13.4 pounds of this product/1,000 sq. ft. per year. In the State of New York do not exceed 0.5 lbs. active ingredient per acre per year (8.8 lbs./1,000 sq. ft.). Use is prohibited in Nassau and Suffolk Counties.		
** May require split or sequential applications for full control.		

Postemergence control is limited; see "Control of Crabgrass" and "Use Precautions" sections above. Regions include areas listed below. See Map of United States.

- North: all other areas.
- Transition: DE, KS, KY, MD, MO, NJ, VA, WV, southeastern PA, southern areas of IL, IN, OH, & Coastal areas of CT, NY, & RI.
- South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX
- Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.



Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the "Program 2" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application may be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control:

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) application spaced 6-8 weeks apart may provide improved weed control.

Control of Other Grass and Broadleaf Weeds

Spring Applications

Used as directed for crabgrass control in the spring, this product will also control (at the "Program 2" rates) the following weeds when applied prior to their emergence:

Barnyardgrass	Lespedeza (common)
Bluegrass (annual)	Oxalis (buttercup, creeping & yellow woodsorrel)
Crowfootgrass	
Foxtail (yellow & green)	Purslane (common)
Goosegrass	Speedwell (corn)
Kikuyugrass	Spurge (prostrat & spotted)
Ryegrass (annual & perennial)	Smutgrass

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 2" rates) the following weeds when applied prior to their emergence:

Bluegrass (annual)	Geranium (Carolina)
Bittercress	Parsley-piert
Chickweed	Pineappleweed
Henbit	Shepherdspurse

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying a uniform distribution of granules.

Spreader (Will list all or some of the following on marketing label)	Spreader Settings Lbs. of product to apply /1,000 sq. ft.				
	2.20 lbs	3.17 lbs	4.41 lbs	6.53 lbs	8.83 lbs
Scotts	X	X	X	X	X
Scotts Drop	X	X	X	X	X
Scotts SpeedyGreen	X	X	X	X	X
Cyclone	X	X	X	X	X
Earthway	X	X	X	X	X
Earthway Broadcast	X	X	X	X	X
Earthway Drop	X	X	X	X	X
Lesco	X	X	X	X	X
Lesco High Wheel	X	X	X	X	X
Spyker	X	X	X	X	X
Prizelawn	X	X	X	X	X
Earthway	X	X	X	X	X

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures.

IF EMPTY: Do not reuse this bag. Place in trash.

IF PARTLY FILLED: Call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to

Pesticide Disposal directions listed above.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container, offer for recycling if available.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase process paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the labels and is reasonably fit for the purposes stated on the labels when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials; manner of application or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount or product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller promptly notifies of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of this Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Distributed by:
Knox Fertilizer Company, Inc.
P.O. Box 248
Knox, Indiana 46534
www.shawsknoxfert.com
Net Weight XX Pounds (XX kg.)

Batch No.

8
11

Dimension® 130 Turf Herbicide with Plant Food

This product is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turf.

Not for use on turf being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

ACTIVE INGREDIENT:

Dithiopyr* 0.13%

INERT INGREDIENTS: 99.87%

Total 100.00%

*CAS # 97886-45-8

Fertilizer Guaranteed Analysis	
Nitrogen	XX%
Phosphate.....	XX%
Potash.....	XX%
Derived from Urea, Ammoniated Phosphate, and Muriate of Potash.	

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA REG. NO. 8378-53

EPA EST. NO. 8378-IN-1

Read the entire label before using this product. Use only according to label instructions.

NOTICE: Before using this product, read the entire Precautionary Statements, Terms and Conditions of Use, Directions for Use, Use Precautions, and Storage and Disposal Instructions. If the Terms and Conditions of Use are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

Net Weight XX lbs. (XX kg.)

-----End Front Panel-----

-----Optional Marketing Statements to add to Front Panel-----

- A Selective Herbicide Plus Fertilizer Combination Product that Provides Control of Crabgrass and Other Weeds in Established Lawns and Ornamental Turfgrasses.
- Premium Granular Fertilizer - Providing the Nutrients Needed for Beautiful Healthy Lawn.
- Contains Dimension® - A Highly Effective Herbicide for Crabgrass Control! Can be Applied Up to 4 Weeks Later Than Other Pre-Emergent Herbicides as an Early Post Emergent Herbicide for Control of Crabgrass.
- This Bag Treats Up to XXX Square Feet.
- Controls Crabgrass and Other Annual Grasses.
- Feeds Your Lawn and Promotes Thick Green Grass.
- Controls Crabgrass.
- Feeds your Lawn.
- XX Nitrogen Slow Release Nitrogen for Continuous Greening.
- XX Phosphorous Helps Develop a Stronger Root System.
- XX Potash Vital for Growth. Improves Plant Hardiness and Drought Resistance.
- Covers XXX Square Feet.

-----End Optional Marketing Statements-----

NOTIFICATION
APR 4 - 2010

9
11

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 from 6:30 a.m. to 4:30 p.m. Pacific Time, everyday, for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS and DOMESTIC ANIMALS
CAUTION: CAUSES EYE IRRITATION. HARMFUL IF INHALED.
 Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS
 This pesticide is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

GENERAL INFORMATION

This product is a combination product that provides both weed control and the required nutrients for optimum turfgrass growth in a single product.

It is a selective herbicide that can be applied up to 4 weeks later than other crabgrass preventers and will provide up to 4 months control of crabgrass.

It will also control other grass and broadleaf weeds listed on this label if applied prior to their seed germination.

This product may be applied to established lawns and ornamental turfgrasses composed of the following tolerant grass species:

- Cool-Season Turfgrasses
 Creeping Bentgrass
 Fines Fescues
 Kentucky Bluegrass
 Perennial Ryegrass
 Tall Fescues

- Warm-Season Turfgrasses
 Bahiagrass
 Bermudagrass
 Buffalograss
 Carpetgrass
 Centipedegrass
 St. Augustinegrass
 Zoysiagrass

This product may be applied in the spring or fall prior to the weed seed germination in established lawns and ornamental turfgrass.

This product provides preemergence control of crabgrass and other weeds when applied prior to their germination, and it may also provide postemergence control of seedling crabgrass. The postemergence control provided by this product is limited to crabgrass. It only will provide control up to 4 weeks after crabgrass germination if applied for post emergent control. Do not apply this product later than 4 weeks after crabgrass germination.

For control of other grasses and broadleaf weeds, this product must be applied as a preventative treatment before their emergence. Only crabgrass will be controlled with applications made 4 weeks after germination.

Improved weed control may occur when this product is applied within a few days after mowing and further mowing is delayed until the turfgrass has been watered or receives adequate rainfall to dissolve the fertilizer granules.

Reseeding, Overseeding, Or Sprigging

Reseeding, overseeding, or sprigging of turfgrasses listed on this label into areas that have been previously treated with this product must be delayed until 8 weeks after application. Reseeding, overseeding, or sprigging before 8 weeks after application may injure or prevent the establishment of the newly reseeded, overseeded or sprigged turfgrass.

Use on New Lawns of Kentucky Bluegrass, Perennial Ryegrass and/or Fall Fescue

To establish a new lawn of Kentucky Bluegrass, Perennial Ryegrass and/or Fall Fescue, this product may be applied 2 weeks after the first sign of germination. For other tolerant turfgrasses listed on this label, wait until the grass is well established (2 mowings) before applying this product.

USE PRECAUTIONS

- This product should only be applied to lawns and ornamental turfgrasses that are well established. Do not apply to lawns or turfgrasses that are under stress.
- DO NOT apply more than 8.8 pounds of this product/ 1,000 sq. ft. per application and no more than 13.4 pounds of this product/ 1,000 sq. ft. per year (1,000 square feet = 100 X 10 feet).
- DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.
- Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

This product will provide control of crabgrass and other grass and broadleaf weeds listed below when applied prior to germination in established lawns and ornamental turfgrasses.

Weeds Controlled by this product:

- Grass Weeds:
 Annual Bluegrass
 Annual Ryegrass
 Barnyardgrass
 Crabgrass
 Crowfootgrass
 Goosegrass
 Green Foxtail
 Kikuyugrass (seedling)
 Perennial Ryegrass
 Smutgrass (seedling)
 Yellow Foxtail

- Broadleaf Weeds:
 Bittercrass
 Buttercup Oxalis
 Carolina Geranium
 Chickweed
 Common Lesperleza
 Common Purslane
 Crn Speedwel
 Creeping Coralis
 Henbit
 Parsley-piert
 Pineappleweed
 Prostrate Spurge
 Spotted Spurge
 Yellow Woodcress

APPLICATION TIMING

This product may be applied either in the spring or in the fall for the control of weeds.

When this product is applied in the fall, a sequential or second application may be made the following spring to provide up to 4 months control of weeds.

If this product is applied in the fall, DO NOT reseed, overseed or sprig turfgrasses before 12 weeks after application of this product due to potential injury to newly reseeded, overseeded or sprigged turfgrasses.

Length of Control:

This product will provide up to 4 months of control of listed weeds when applied according to label directions.


The actual length of control will vary depending on use rate, weather conditions, conditions of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States that this product is being applied.

Please check the map below to determine the region of the U.S. that applies to your particular use area, and find the number of pounds of this product needed in your area based on the length of weed controlled given in the tables below.

Post Emergence Crabgrass Control

For post emergent crabgrass control, apply this product at the recommended rates when crabgrass is small (3-4 leaf). This product will not control large tillered crabgrass. Applications will control existing crabgrass and prevent further emergence.

Use "Regions" within the United States



RECOMMENDED USE RATES (Pounds of this product to apply 1,000 square feet of turfgrass)			
Length of Weed Control			
Regions of U.S.	2-3 Months Control	3-4 Months Control	Early Post Emergence Control
North	2.2 (0.13 lb. AI/acre)	3.1 (0.17 lb. AI/acre)	4.0- 8.8 (0.22-0.50 lb. AI/acre)*
South	3.4 (0.19 lb. AI/acre)	4.0 - 4.5 (0.22-0.26 lb. AI/acre)*	4.0- 8.8 (0.22-0.50 lb. AI/acre)*

DO NOT apply more than 8.8 pounds of this product per 1,000 sq. ft. per application and no more than 13.4 pounds of this product/ 1,000 sq. ft. per year (1,000 square feet = 100 X 10 feet).

In the State of New York: Do not exceed 0.25 pounds active ingredient Dithiopyr per acre per year (4.4 lbs. product/1000 sq. ft.). Do Not Apply More Than Once Per Year To The Same Area.

*Use the higher rate in areas of high weed pressure.

APPLICATION EQUIPMENT AND INSTRUCTIONS

This product can be applied with a drop or rotary spreader designed to apply granular herbicides.

For best results, apply this product evenly and uniformly over the turfgrass area avoiding streaking, skips or overlaps. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands.

Before each application, calibrate the spreader according to the equipment manufacturer's directions. Adjust the spreader settings such that the spreader delivers the appropriate application rates of this product as recommended above.

A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then applying the remaining one-half of this product at a right angle to the previous application direction.

Check application equipment frequently to insure equipment is functioning properly and applying a uniform distribution of this product.

Spreader Settings
Lbs. of product to apply/1,000 sq. ft.

Spreader (Will list all or some of the following on marketing label)	2.2 lbs.	3.1 lbs.	3.4 lbs.	4.0 lbs.	4.5 lbs.	8.8 lbs.
Cyclone	X	X	X	X	X	X
Scotts	X	X	X	X	X	X
Earthway	X	X	X	X	X	X
Earthway Best	X	X	X	X	X	X
Earthway Drop	X	X	X	X	X	X
Lesco	X	X	X	X	X	X
Lesco High Wheel	X	X	X	X	X	X
Prizelawn	X	X	X	X	X	X
Ortho Drop (5000)	X	X	X	X	X	X
Ortho Rotary (6000)	X	X	X	X	X	X
Precision Red Devil Rotary	X	X	X	X	X	X
Prizelawn	X	X	X	X	X	X
Republic EZ Drop	X	X	X	X	X	X
Republic EZ Grow	X	X	X	X	X	X
Scotts Drop	X	X	X	X	X	X
Scotts AccuGreen	X	X	X	X	X	X
Scotts Easy Green	X	X	X	X	X	X
Scotts SpeedyGreen	X	X	X	X	X	X
Spyker	X	X	X	X	X	X

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PESTICIDE DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Do not reuse this bag. Place in trash.

IF PARTLY FILLED: Call your local solid waste agency for disposal.

11
11

Instructions: Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to Pesticide Disposal directions listed above.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container, offer for recycling if available.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid.

Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the labels and is reasonably fit for the purposes stated on the labels when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, manner of application or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount or product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller promptly notifies of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of this Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or excess the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Manufactured by:
Knox Fertilizer Company, Inc.
PO Box 248
Knox, Indiana 46534

Net Weight XX lbs. (XX kg.)

Batch No.