

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	EPA REGISTRATION NO. 8378-56	DATE OF ISSUANCE FEB 6 1998
	TERM OF ISSUANCE Conditional	
	NAME OF PESTICIDE PRODUCT SHAW'S TURF FOOD/CRABGRASS CONTROL BARRICADE® 50	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Knox Fertilizer Company
 P.O. Box 248
 Knox, IN 46534

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Add the phrase "EPA Registration No. 8378-56" to the label before you release the product for shipment.
2. Delete the phrase, "(83E50amd.265/G1-28-98)" from the first panel statements and similar notations on second page labeling.
3. Delete all repetitious and redundant information of second page of labeling. Only the described use-patterns for "Landscape Ornamentals Including Established Perennials and Wildflower Plantings" from that page will remain. The redundant information includes main headings, sub-headings, ingredient statements, precautionary labeling down to "Application Directions", except for the second heading on that page.
4. Submit one (1) copy of the final printed labeling described above before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Eugene M. Wilson
 Eugene M. Wilson
 Acting Product Manager (23)
 Herbicide Branch
 Registration Division (7505C)

Enclosure
 ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------

Shaw's Turf Food / Crabgrass Control Barricade® 50

(20 lbs. treats up to 6,710 sq.ft.)

For Selective Preemergence Control of Grass and Broadleaf Weeds in Established Turf Grasses, (excluding golf course putting greens) and Lawns - Also For Landscape Ornamentals Including Established Perennials and Wildflower Plantings.


ACTIVE INGREDIENT:
Prodiamine [N,N-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] - 0.50%
INERT INGREDIENTS: - - - - 99.50%
Total 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See the following section, Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information. Barricade® is a Registered Trademark of Novartis.

EPA Reg. No. 8378-XX EPA Est. No. 8378-in-1

Crabgrass Seed Germination Dates



Approximate Date

1. Mid-north & northwestern After May 30
2. Upper/mid-states & northeast After May 10
3. Lower mid-states After April 20
4. Upper southern states After Mar 20
5. Warm season states Jan 1 to Mar 20

USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate/Calendar Year of Product by Turf Grass Species⁽¹⁾

Turf Species	Lbs. Product/A	Lbs. a.l./A
Creeping Bentgrass	130 lbs.	0.65 lb.
Creeping Red Fescue	150 lbs.	0.75 lb.
Buffalograss	200 lbs.	1.00 lb.
Kentucky Bluegrass	"	"
Perennial Ryegrass	"	"
Bermudagrass ⁽²⁾	300 lbs.	1.50 lbs.
Bahia grass	"	"
Centipede grass	"	"
Seashore Paspalum	"	"
St. Augustine grass	"	"
Tall Fescue	"	"
Including turf type Zoysia	" (1A = per Acre)	"

⁽¹⁾These are the maximum rates per calendar year by species limitations. ⁽²⁾May be used on newly-sprigged or plugged Bermudagrass at rates not to exceed 104 lbs./A. Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations. Do not apply more than 300 lbs. of this product per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates section.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

KNOX FERTILIZER CO.

P.O. Box 248 - Knox, IN 46534 U.S.A.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds.

Barnyardgrass	Herb	Purslane, common
Bluegrass, Annual (Poa Annu)	Grass	Pusley, Florida
Carpelweed	Johnsongrass (from seed)	Rescuegrass ⁽¹⁾
Chickweed, Common	Juncelence	Shepherd's Purse ⁽¹⁾
Chickweed, Mouseear	Knockweed	Signalgrass, Briff.
(from seed)	Kochia	Speedwell, Persian
Crabgrass (Large, Smooth)	Lambsquarter, common	Springletop
Crowfootgrass	Lovgrass	Surge, Prostrate
Cupgrass, Woolly	Panicum (Texas, Fall, Brownlop)	Witchgrass
Foxtails, Annual	Pigweed	Woodsorrel, Yellow
Goosegrass ⁽¹⁾		(from seed)

For weed reference numbers⁽¹⁾ refer to the following sentences.

(1) In many areas a single application of 130 to 300 lbs./A will control goosegrass. However under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 130 to 200 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rates section. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rates section. (2) Application for this weed should be made in late summer, fall, or winter prior to germination. (3) Suppression only.

Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

SPREADER SETTINGS FOR 20 LBS. 6,710 SQ.FT. 5,815 SQ.FT. 4,360 SQ.FT. 2,900 SQ.FT.				
Scott's Evergreen 18*	# 5.0	# 5.5	# 6.0	# 7.0
Scott's Speedy Brdct	# 3.0	# 3.5	# 4.0	# 4.5
Red Devil Broadcast	# 5.0	# 5.5	# 6.0	# 7.0
Republic EZ Drop 18*	# 6.0	# 7.0	# 8.0	# 8.5
Republic EZ Broadcast	# 2.5	# 3.0	# 4.0	# 5.0
Republic EZ Trtr. Brdct	# 3.5	# 4.5	# 5.5	# 6.0

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Product Rate lbs./Acr	product	a.l./A	Months Before Overseeding		
			North	Transition	South
100	0.50		4	4	4
130	0.65		5	4	4
150	0.75		6	5	5
160	0.80		-	6	6
200	1.00		-	7	7
228	1.14		-	-	9
260	1.30		-	-	10
300	1.50		-	-	12

GUARANTEED FERTILIZER ANALYSIS
Total Nitrogen (N) variable %
Available Phosphoric Acid (P₂O₅) variable %
Soluble Potash (K₂O) variable %
Ingredients: Ureaformaldehyde, Urea, Sulfur Coated Urea, Ammonium Phosphate, Muriate of Potash



**ACCEPTED
with COMMENTS
In EPA Letter Dated
FEB 6 1988**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

8378-56

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

Statements of Practical Treatment

Eyes: Flush thoroughly with water for several minutes. **Medical attention** if irritation persists. **On Skin:** Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists. **If Inhaled:** Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters. If product is spilled, sweep up and use per label instructions or dispose of as indicated in the STORAGE AND DISPOSAL section.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Do not reuse empty bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or at an incineration facility, or if allowed by State and local authorities, by burning locally. If burned, stay out of smoke.

DIRECTIONS FOR USE

Violation of Federal law to use this product in a manner inconsistent with its labeling. This product is a turf fertilizer and a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), and lawns, landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least by 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown: (1) for sale or other commercial use; (2) for commercial seed production or; (3) for research purposes.

For use on plants intended for esthetic purposes or climatic modification and being grown in ornamental gardens, or parks, or on golf courses, or lawns and grounds.

Do not graze or feed livestock forage out from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

Application Directions

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates. (83B50amd.P6501-28-98)

Net Weight 17.25;20;23;26;34.5;40;46;50 Pounds

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

Statements of Practical Treatment

In Eyes: Flush thoroughly with water for several minutes. medical attention if irritation persists. **On Skin:** Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists. **Inhaled:** Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters. If product is spilled, sweep up and use per label instructions or dispose of as indicated in STORAGE AND DISPOSAL section.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container away from other fertilizer, feed or foodstuffs, and separated from other pesticides. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Do not reuse empty bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or at an incineration facility, or if allowed by State and local authorities, by burning locally, if burned stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in landscape ornamentals, established perennials and wildflower plantings. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown: (1) for sale or other commercial use; (2) for commercial seed production or; (3) for research. For use on plants intended for esthetic purposes or climatic modification and being grown in ornamental gardens, or parks, or on golf courses, and grounds. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aurally. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

Application Directions

Apply uniformly through suitable, calibrated, hand held, push or pull type granule spreader application equipment.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weed. The length of time of residual weed control provided by this product is related to the rate applied.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

- | | | |
|---------------------------------|--------------------------|--------------------------------|
| Barnyardgrass | Henbit | Purslane, common |
| Bluegrass, Annual (For Annual) | Knightgrass | Pusley, Florida |
| Carpetweed | Johnsongrass (from seed) | Rescuegrass ¹⁾ |
| Chickweed, Common | Junglecica | Shepherd's Purse ²⁾ |
| Chickweed, Mouseear (from seed) | Knotweed | Signalgrass, Brill. |
| | Kochia | Speedwell, Persian |
| Crabgrass (Large, Smooth) | Lambquarter, common | Spangletop |
| Crowfootgrass | Lovegrass | Surge, Prostrate |
| Cupgrass, Woolly | Panicum (Texas, Fall) | Witchgrass |
| Foxtails, Annual | Browntop | Woodsorrel, Yellow |
| Goosegrass ¹⁾ | Pigweed | (from seed) |

For weed reference numbers¹⁾ refer to the following sentences.

(1) In many areas a single application of 130 to 300 lbs./A will control goosegrass. However under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 130 to 200 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed a total of 300 lbs. per season. (2) Application for this weed should be made in late summer, fall, or winter prior to germination (3) Suppression Only. Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rate of a total of 300 lbs. All applications must be made prior to germination of the weed seeds. (bar50-or-P65(01-29-98))

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	variable %
Available Phosphoric Acid (P ₂ O ₅)	variable %
Soluble Potash (K ₂ O)	variable %
Ingredients: Ureaformaldehyde, Urea, Sulfur Coated Urea, Ammonium Phosphate, Muriate of Potash	

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

Net Weight (5),(10),(20),(40),(60) Pounds

Knox Fertilizer Co. Inc.
P.O. Box 248 - Knox, IN 46534 U.S.A.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product into the soil.

Newly Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name	Common Name
<i>Abies</i> spp.	Fir species* (Balsam, Fraser, Noble, etc.)
<i>Abelia grandiflora</i>	Abelia; Sherwood
<i>Acer palmatum</i>	Japanese Maple
<i>A. platanoides</i>	Norway Maple
<i>Acillilea</i> , sp.	Yarrow; King Edward
<i>Actinidia chinensis</i>	Kiwi
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus orientalis</i>	
<i>Akalia quintata</i>	Five-Leaf or Chocolate Vine
<i>Allium cernuum</i>	Lady's Leak, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia</i> sp.	Aquilegia; Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctostaphylos calandula</i>	Cape Weed
<i>Artemisia</i> sp.	Wormwood; Silver Mound, Castle Aster
<i>Aster</i> sp.	Aster; Bonny Blue, Purple Dome
<i>Aster X Frikartii</i>	Lady Fern; Fern Lady
<i>Athrium Filiz-femina</i>	Japanese Aucuba
<i>Aucuba japonica</i>	Fibrous Begonia; Hardy Grandis
<i>Begonia</i> sp.	Barberry
<i>Berberis gladiwomensis</i>	Wintergreen Barberry
<i>B. julianae</i>	Mentor Barberry
<i>B. mentorensis</i>	Japanese Barberry
<i>B. thunbergii</i>	Warty Barberry
<i>B. verruculosa</i>	
<i>Borgeria cordifolia</i>	Snowbank
<i>Boitonia asteroides</i>	Bougainvillea
<i>Bougainvillea</i> spp.	Butterfly-Bush (Dwarf Blue); Royal Red
<i>Buddleia davidii</i>	Japanese Boxwood
<i>Buxus microphylla</i>	Weeping Bottlebrush
<i>Callistemon viminalis</i>	Scotch Heather
<i>Calluna vulgaris</i>	Tussock Bellflower (White Clips)
<i>Campanula carpatica</i>	Trumpet Creeper, Trumpet Flower, Madame Galen
<i>Campis X Tagilabuana</i>	Hottentot Fig (Ice Plant)
<i>Carpobrotus edulis</i>	Feathery Cassia
<i>Cassia artemisoides</i>	Wild Lilac
<i>Ceanothus rigidus</i>	
<i>Ceratostigma plumbagonoides</i>	False Cypress
<i>Chamaecyparis pisifera</i>	
<i>Chrysanthemum nipponicum</i>	Cleyera
<i>Cleyera japonica</i>	Citrus species*
<i>Citrus</i> spp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
<i>Coreopsis</i> sp.	Flowering Dogwood
<i>Cornus florida</i>	American Dogwood
<i>C. stolonifera</i>	Pampas Grass
<i>Cortaderia selkiana</i>	Crabber Cotoneaster
<i>Cotoneaster apiculatus</i>	Cotoneaster
<i>C. buxifolia</i>	Cotoneaster
<i>C. dammeri</i>	Bearberry Cotoneaster
<i>C. macrophylla</i>	Rockspray Cotoneaster
<i>Cretaegus</i> spp.	Hawthorne
<i>Crocosmia</i> sp.	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma</i> sp.	Cooper Pink
<i>Delphinium</i> sp.	Larkspur; Blue Elf
<i>Dianthus deltoides</i>	Dianthus; Maiden Pink; "Zing"
<i>D. granatopolitanus</i>	Cheddar Pink
<i>Dodonaea viscosa</i>	Hop Bush
<i>Echinacea purpurea</i>	Coneflower, Purple; Magnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>E. japonica</i>	Japanese Spindle Tree (Evergreen Euonymus)
<i>E. klatschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>F. suspensa</i>	Weeping Forsythia
<i>F. viridissima</i>	Greenstem Forsythia
<i>Gaillardia</i> sp.	Gaillardia, Blanket Flower; "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-jasmine, Radicans
<i>Gaura</i> sp.	
<i>Gentiana dahurica</i>	Gentian
<i>Gentianum cinereum</i>	Cranesbill
<i>Gladiolus</i> spp.	Gladiolus species**
<i>Gypsophila repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellianthomum</i> sp.	Sunrose
<i>Hemerocallis</i> sp.	Daylily; Aztec Gold, Stella De Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus</i> sp.	Mallow; Disco Belle White
<i>Hibiscus rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. sieboldiana</i>	Hosta, "Searsucker"
<i>Houttuynia cordata</i> var. <i>variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea

<i>L. japonicum</i>	<i>Liriope muscari</i>	<i>Liriope muscari</i> var. <i>variegata</i>	<i>Liriope</i> , Creeping
<i>L. lucidum</i>	<i>Lobelia cardinalis</i>	<i>Lobelia spicata</i>	Cardinal Flower, Indian Pink
<i>Lilium</i> sp.	<i>Lonkera japonica</i>	<i>Lobelia cardinalis</i>	Japanese Honeysuckle
<i>Liriope muscari</i>	<i>L. tatarica</i>	<i>Loropetalum chinense</i>	Tatarian Honeysuckle
<i>Liriope muscari</i> var. <i>variegata</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Burgundy
<i>Liriope spicata</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Loosestrife; Modern Pink
<i>Lobelia cardinalis</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Magnolia species**
<i>Lonkera japonica</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Ice Plant
<i>L. tatarica</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Crabapple*
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Yaku Jima, Silberfeder
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Heavenly Bamboo
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Narcissus species**
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Oleander
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Oliver
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Mondo Grass
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Osmanthus (False Holly); Gulf Tide
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Trailing African Daisy
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Sourwood
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Tree Peony
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Fountain Grass (Dwarf)
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Avocado*
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Fraser's Photinia (Redtip)
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Dragonhead, False; Vivid Spruce species** (Colorado Blue, Norway, etc.)
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Lily-of-the-Valley Bush
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Calabrian Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Canary Island Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Slash Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Aleppo Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Austrian Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Longleaf Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Monterey Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Japanese Black Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Eastern White Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Scotch Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Lablolly Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Virginia Pine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Pisachio*
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Queensland Pittosporum
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Japanese Yew
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	English Laurel
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Daucus
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Firethorn (Scarlet)
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Firethorn
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Bradford Pear sp.
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Oak, Shumard's Red
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Oak species
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Indian Hawthorne
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Yedda Hawthorn
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Coral Bells"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Delaware Valley White"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Flame Creeper"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Formosa"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Girard Crimson"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"George L. Tabor"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Hino-crimson"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"J.M."
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Rosaum Elegans"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"Wakefields"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	"White Gump"
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Lady Bank's Rose
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Rosemary*
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Black-Eyed Susan; Goldstrum
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Santolina
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Saxifrage; Purple Dome
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Pincushion Flower
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Stonecrop
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Stonecrop; Lidakense
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Stonecrop
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Stonecrop; Dragon's Blood
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Spiraea; Anthony Waterer
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Australian Brushcherry
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Japanese Yew
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Yew
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Cermander
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Meadow Rue
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	American Arborvitae
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Star Jasmine
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Canada Hemlock
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Tulip species
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Veronica, Speedwell; Sunny
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Border
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Japanese Viburnum
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Sweet Viburnum
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Japanese Snowball
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Canary Island Viburnum
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Arrowood Viburnum
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Launstinus
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Cranberry Bush
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Leatherleaf Viburnum
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Greater Periwinkle, Vinca
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Common Periwinkle, Vinca
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Grape*
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Old Fashioned Weigela
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Spanish Bayonet, Yucca
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>	Adam's Needle, Yucca
<i>Loropetalum chinense</i>	<i>Loropetalum chinense</i>		