

FRONT PANEL

DISINFECTANT SOLUTION (PUMP SPRAY)

Contamination and Odor Control

SPORICIDIN
Brand
Disinfectant Solution

Continuous Residual Activity
for Over 6 Months*

ACTIVE INGREDIENTS:

Phenol 1.56%
Sodium Phenate 0.06%

- BACTERICIDAL
- FUNGICIDAL
- VIRUCIDAL**
- TUBERCULOCIDAL
- ELIMINATES ODORS*
- PREVENTS MOLD AND MILDEW

INERT INGREDIENTS: 98.38%

Solution is alkaline, pH 7.9-8.6


Keep Out of Reach of Children

CAUTION:

See additional precautions on back panel

EPA Reg. No. 8383-3

Net Wt. 22 fl. oz. (651 ml)



HOSPITAL DISINFECTION

An effective disinfectant and active deodorant for hospitals, clinics, medical and veterinary offices, laboratories, industrial clean rooms, nursing homes, ambulances, hotels, restaurants, schools, airplanes, trains, boats, autos, buses, health spas and toilets.

DISINFECTANT. Acts effectively on germ-laden non-porous equipment and surfaces in medical treatment and patient rooms, operating rooms, emergency rooms, recovery rooms and bathrooms. For use on medical equipment and beds, surgical carts, counter tops, manikins, telephones, furniture, wheelchairs, walkers, sinks, floors, walls, light switches, linen hampers, shower stalls, toilet seats, trash containers, damp and musty areas, including animal enclosures. Reduces the hazards of cross-contamination on environmental surfaces.

ACTIVE DEODORIZER. *It provides continuous residual bacteriostatic activity against odor-causing organisms, *for over 6 months* in the presence of adequate moisture if not removed from surfaces including plastics, latex, vinyl, glass, wood, metal, porcelain, tile and paint.

♦ Kills Athlete's Foot fungus on bathroom, shower, and pool surfaces. ♦ Self-sanitizes surfaces against odor-causing bacteria. ♦ Penetrates cracks. ♦ Non-staining ♦ Contains no harmful abrasives.

SPORICIDIN Brand DISINFECTANT SOLUTION is germicidal when used as directed for disinfection but differs in composition from other Sporcidin products.

PASSES THE AOAC EFFICACY STANDARDS FOR INSTITUTIONAL AND HOSPITAL DISINFECTION. Tested against *Staphylococcus aureus*, *Streptococcus pyogenes*, *Streptococcus salivarius*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, *Escherichia coli*. **Herpes simplex types 1 and 2 (oral and genital), Influenza A₂ (Japan 305/57 Asian strain), Vaccinia, and Polio type 1 viruses. HIV-1 (AIDS virus) on environmental surfaces, Trichophyton mentagrophytes (Athlete's Foot fungus) and *Mycobacterium tuberculosis* on environmental surfaces.

DIRECTIONS:

It is a violation of U.S. Federal Law to use this product in a manner inconsistent with its labeling.

PRECLEAN surface to remove soil and filth. Wipe dry with paper towel, cloth or sponge.

TO DISINFECT AND DEODORIZE: Thoroughly wet pre-cleaned surface with disinfectant. Ensure surface remains moist for a 10 minute contact time at room temperature (20° C/68° F or above). **NOTE:** See Sporcidin Bulletin No. 303 for HIV-1 (AIDS virus) instructions.

TO PREVENT MOLD AND MILDEW: Repeat application weekly, or sooner if new growth appears.

FOR FOGGING CLEAN ROOMS: See Sporcidin Bulletin No. 301.

CAUTION: Avoid direct contact with food. In case of eye contact, immediately flush with water. If necessary, consult a physician.

| |
|---|
| STORAGE AND DISPOSAL: Store at room temperature. Discard empty container in regular trash. |
|---|

SPORICIDIN BULLETIN NO. 301
(EPA Reg. No. 8383-3)

FOGGING TREATMENT FOR ARTICLES AND SURFACES IN SEALED ROOMS AND CUBICLES

Directions:

It is a violation of U.S. Federal Law to use this product in a manner inconsistent with its labeling.

Sporicidin Brand Disinfectant Spray is recommended for fogging articles and surfaces in sealed rooms and cubicles as an adjunct to manual cleaning and disinfecting procedures. Unscrew pump bottle and charge the fogging device with the required volume of solution. Seal all windows and doors. Set timer for treatment cycle as recommended by device manufacturer. Turn on device, leave room and reseal door. Do not allow anyone to re-enter room or cubicle for at least 2 hours after fogging cycle has been completed. Upon re-entering room or cubicle, follow regular cleaning and disinfecting procedures.

Precautions:

Avoid skin contact: Possibility of sensitization exists and may cause skin irritation. Flush thoroughly with water after contact.

Avoid eye contact: Causes eye irritation. In case of contact, flush with water immediately and get medical attention.

Harmful if swallowed. Drink large quantities of water and call physician immediately.

Avoid food contamination.

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER. DISPOSE OF IN PROPER RECEPTACLE.

9/88

SPORICIDIN BULLETIN NO. 303:
Sporicidin Brand
Disinfectant Solution and Towellette
(EPA Reg. Nos. 8383-3 and 8383-7)

- a. **Personal Protection:**
Persons handling items soiled with blood or body fluids likely to be contaminated with HIV-1 (AIDS virus) must protect themselves by use of disposable latex gloves, gowns, masks and eye covering.
- b. **Decontamination of Contaminated Objects:**
Immediately after use, items contaminated with blood, body fluids, or other soils that may contain HIV-1 (AIDS virus) must be thoroughly wetted with Sporicidin brand disinfectant solution for at least 1 minute before further processing.
- c. **Cleaning Procedure:**
Following decontamination, objects and surfaces are moved to the cleaning and preparation areas where they are washed, rinsed and rough dried to remove all blood and body fluids before application of the disinfectant.
- d. **Disinfection:**
Thoroughly wet entire object or surface with solution or towellette. HIV-1 (AIDS virus) is killed within 1 minute contact time. *Note: Use a 10 minute contact time for other organisms.
- e. **Disposal of Infectious Materials:**
Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

Sporicidin

BULLETIN NOS.
303, 403 and 503

**SPECIAL INSTRUCTIONS
FOR
CLEANING AND
DECONTAMINATION OF
SURFACES/OBJECTS
SOILED WITH
BLOOD/BODY FLUIDS
AGAINST
HIV-1 (AIDS VIRUS)**