FRONT PANEL

DISINFECTANT SOLUTION (PUMP SPRAY)

Contamination and Odor Control

SPORICIDIN Brand Disinfectant Solution

Continuous Residual Activity for Over 6 Months*

ACTIVE INGREDIENTS:

Phenol 1.56% Sodium Phenate 0.06%

INERT INGREDIENTS: 98.38%

Solution is alkaline, ph 7.9-8.6

- BACTERICIDAL
- FUNGICIDAL
- VIRUCIDAL**
- TUBERCULOCIDAL
- ELIMINATES ODORS*
- PREVENTS MOLD AND MILDEW


Keep Out of Reach of Children

CAUTION:

See additional precautions or back panel

EPA Reg. No. 8383-3

Net Wt. 22 fl. oz. (651 ml)



VESTADO LA E COPY

HOSPITAL DISINFECTION

An effective disinfectant and active deodorant for hospitals, clinics, medical and veterinary offices, laboratories, industrial clean rooms, homes, nursing homes, ambulances, hotels, restaurants, schools, airplanes, trains, boats, autos, buses, health spas and toilets.

DISINFECTANT. Acts effectively on germ-laden non-porous equipment and surfaces in medical treatment and patient rooms, operating rooms, emergency rooms, recovery rooms and bathrooms. For use on medical equipment *** and beds, surgical carts, counter tops, manikins, telephones, furniture, wheelchairs, walkers, sinks, floors, walls, light switches, linen hampers, shower stalls, toilet seats, trash containers, damp and musty areas, including animal enclosures. Reduces the hazards of cross-contamination on environmental surfaces.

ACTIVE DEODORIZER. *It provides continuous residual bacteriostatic activity against odor-causing organisms, for over 6 months in the presence of adequate moisture if not removed from surfaces including plastics, latex, vinyl, glass, wood, metal, porcelain, tile and paint.

*Kills Athlete's Foot fungus on bathroom, shower, and pool surfaces. *Self-sanitizes surfaces against odor-causing bacteria. *Penetrates cracks. *Non-staining. *Contains no harmful abrasives.

ORICIDIN Brand DISINFECTANT SOLUTION is germicidal when used as directed for disinfection but ifers in composition from other Sporicidin products.

PASSES THE AOAC EFFICACY STANDARDS FOR INSTITUTIONAL AND HOSPITAL DISINFECTION. Tested against Staphylococcus aureus, Streptococcus pyogenes, Streptococcus salivarius, Salmonella choleraesuis, Pseudomonas aeruginosa, Escherichia coli, **Herpes simplex types l and 2 (oral and genital), Influenza A₂ (Japan 305/57 Asian strain), Vaccinia, and Polio type l viruses. HIV-1 (AIDS virus) on environmental surfaces, Trichophyton mentagrophytes (Athlete's Foot fungus) and Mycobacterium tuberculosis on environmental surfaces.

DIRECTIONS:

It is a violation of U.S. Federal Law to use this product in a manner inconsistent with its labeling.

PRECLEAN surface to remove soil and filth. Wipe dry with paper towel, cloth or sponge.

TO DISINFECT AND DEODORIZE: Thoroughly wet pre-cleaned surface with disinfectant. Ensure surface remains moist for a 10 minute contact time at room temperature (20°C/68°F or above). NOTE: See Sporicidin Bulletin No. 303 for HIV-1 (AIDS virus) instructions.

TO PREVENT MOLD AND MILDEW: Repeat application weekly, or sooner if new growth appears.

FOR FOGGING CLEAN ROOMS: See Sporicidin Bulletin No. 301.

TO CLEAN AIR DUCTS: Manually apply to hard non-porous precleaned duct surfaces. Rinse surfaces after application.

CAUTION: Avoid direct contact with food. In case of eye contact, immediately flush with water. If necessary, consult a physician.

STORAGE AND DISPOSAL; Store at room temperature. Discard empty container in regular trash.

*** Other than critical and semi-critical items.

SPORICIDIN INTERNATIONAL

5901 Montrose Road Rockville, ND 20852

8383-3

17 NOV 1794

BEST AVAILABLE COPY