

PM 32

9383-4

10/2

FRONT PANEL

DISINFECTANT (AEROSOL) SPRAY

Contamination and Odor Control

SPORICIDIN
Brand
Disinfectant Spray

Continuous Residual Activity
for Over 6 Months*

ACTIVE INGREDIENTS:

Phenol 1.56%
Sodium Phenate 0.06%

INERT INGREDIENTS: 98.38%

Solution is alkaline, pH 7.9-8.6

- BACTERICIDAL
- FUNGICIDAL
- VIRUCIDAL**
- TUBERCULOCIDAL
- ELIMINATES ODORS*
- PREVENTS MOLD AND MILDEW


Keep Out of Reach of Children

WARNING:

See additional precautions on back panel

EPA Reg. No. 8383-4

Net Wt. 22 oz.



2012

BACK PANEL

DISINFECTANT (AEROSOL) SPRAY

HOSPITAL DISINFECTION

An effective disinfectant and active deodorant for hospitals, clinics, medical and veterinary offices, laboratories, industrial clean rooms, homes, nursing homes, ambulances, hotels, restaurants, schools, airplanes, trains, boats, autos, buses, health spas and toilets.

DISINFECTANT. Acts effectively on germ-laden non-porous equipment and surfaces in medical treatment and patient rooms, operating rooms, emergency rooms, recovery rooms and bathrooms. For use on medical equipment *** and beds, surgical carts, counter tops, manikins, telephones, furniture, wheelchairs, walkers, sinks, floors, walls, light switches, linen hampers, shower stalls, toilet seats, trash containers, damp and musty areas, including animal enclosures. Reduces the hazards of cross-contamination on environmental surfaces.

ACTIVE DEODORIZER. *It provides continuous residual bacteriostatic activity against odor-causing organisms, for over 6 months in the presence of adequate moisture if not removed from surfaces including plastics, latex, vinyl, glass, wood, metal, porcelain, tile and paint.

◆Kills Athlete's Foot fungus on bathroom, shower, and pool surfaces. ◆Self-sanitizes surfaces against odor-causing bacteria. ◆Penetrates cracks. ◆Non-staining. ◆Contains no harmful abrasives.

SPORICIDIN Brand DISINFECTANT SPRAY is germicidal when used as directed for disinfection but differs in composition from other Sporicidin products.

MEETS THE AOAC EFFICACY STANDARDS FOR INSTITUTIONAL AND HOSPITAL DISINFECTION. Tested against Staphylococcus aureus, Streptococcus pyogenes, Streptococcus salivarius, Salmonella choleraesuis, Pseudomonas aeruginosa, Escherichia coli, **Herpes simplex types 1 and 2 (oral and genital), Influenza A, (Japan 305/57 Asian strain), Vaccinia, and Polio type 1 viruses. HIV-1 (AIDS virus) on environmental surfaces, Trichophyton mentagrophytes (Athlete's Foot fungus) and Mycobacterium tuberculosis on environmental surfaces.

DIRECTIONS:

It is a violation of U.S. Federal Law to use this product in a manner inconsistent with its labeling.

PRECLEAN surface to remove soil and filth. Wipe dry with paper towel, cloth or sponge.

TO DISINFECT AND DEODORIZE: Shake well before using. Hold spray nozzle 6" to 8" from surface. Thoroughly wet pre-cleaned surface with disinfectant. Ensure surface remains moist for a 10 minute contact time at room temperature (20°C/68°F or above). NOTE: See Sporicidin Bulletin No. 403 for HIV-1 (AIDS virus) instructions.

TO PREVENT MOLD AND MILDEW: Repeat application weekly, or sooner if new growth appears.

ELIMINATE ODORS; Spray 2-4 seconds toward center of room.

TO CLEAN AIR DUCTS: Manually apply to hard non-porous precleaned duct surfaces. Rinse surfaces after application.


WARNING: Avoid direct contact with food. In case of eye contact, immediately flush with water. If necessary, consult a physician.

STORAGE AND DISPOSAL; Store at room temperature. Discard empty container in regular trash.

CONTENTS UNDER PRESSURE: Do not use near fire, sparks, or flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

*** Other than critical and semi-critical items.

SPORICIDIN INTERNATIONAL



5901 Montrose Road
Rockville, MD 20852