

Reg # 8383-7

PM-32

193

FRONT PANEL

DISINFECTANT TOWELETTE

Contamination and Odor Control

SPORICIDIN
Brand
Disinfectant Towelette

Continuous Residual Activity
for Over 6 Months*

ACTIVE INGREDIENTS:

Phenol 1.56%
Sodium Phenate 0.06%

- BACTERICIDAL
- FUNGICIDAL
- VIRUCIDAL**
- TUBERCULOCIDAL
- ELIMINATES ODORS*
- PREVENTS MOLD AND MILDEW

INERT INGREDIENTS: 98.38%

Solution is alkaline, pH 7.9-8.6

Keep Out of Reach of Children

CAUTION:


Avoid direct food contact. On eye contact, flush with water.
If necessary, contact physician.

Contents:

(1) Towelette saturated
with disinfectant.

EPA Reg. No. 8383-7

BEST AVAILABLE COPY



HOSPITAL DISINFECTION

293

An effective disinfectant and active deodorant for hospitals, clinics, medical and veterinary offices, laboratories, industrial clean rooms, nursing homes, ambulances, hotels, restaurants, schools, airplanes, trains, boats, autos, buses, health spas and toilets.

DISINFECTANT. Acts effectively on germ-laden non-porous equipment and surfaces in medical treatment and patient rooms, operating rooms, emergency rooms, recovery rooms and bathrooms. For use on medical equipment*** and beds, surgical carts, counter tops, manikins, telephones, furniture, wheelchairs, walkers, sinks, floors, walls, light switches, linen hampers, shower stalls, toilet seats, trash containers, damp and musty areas, including animal enclosures. Reduces the hazards of cross-contamination on environmental surfaces.

DEODORANT. *It provides continuous residual bacteriostatic activity against odor-causing organisms, for over 6 months in the presence of adequate moisture if not removed from surfaces including vinyl, glass and wood.

♦ Non-staining

♦ Contains no harmful abrasives

SPORICIDIN Brand DISINFECTANT TOWELETTES are germicidal when used as directed but differ in composition from other Sporcidin products.

PASSES THE AOAC EFFICACY STANDARDS FOR INSTITUTIONAL AND HOSPITAL DISINFECTION such as *Staphylococcus aureus*, *Streptococcus pyogenes*, *Streptococcus salivarius*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, *Escherichia coli*. **Herpes simplex types 1 and 2 (oral and genital). Influenza A₂ (Japan 305/57 Asian strain), Polio type 1 virus, HIV-1 (AIDS virus) on environmental surfaces, *Trichophyton mentagrophytes* (Athlete's Foot fungus) and *Mycobacterium tuberculosis* on environmental surfaces.

DIRECTIONS:

It is a violation of U.S. Federal Law to use this product in a manner inconsistent with its labeling.

PRECLEAN surface to remove soil and filth. Wipe dry.

TO DISINFECT AND DEODORIZE: Thoroughly wet surface with towelette. Ensure surface remains moist for a 10 minute contact time at room temperature (20°C/68°F or above).

TO CLEAN AIR DUCTS: Manually apply to hard non-porous precleaned duct surfaces. Rinse surfaces after application.

TO PREVENT MOLD AND MILDEW: Repeat application weekly or sooner if new growth appears.

NOTE: See Sporcidin Bulletin No. 303 for HIV-1 (AIDS virus) instructions.

STORAGE AND DISPOSAL


STORAGE: Store at room temperature.

DISPOSAL: Discard used towelettes and empty container in regular trash.


*** Other than critical and semi-critical items.

Sporcidin
International

5901 Montrose Road
Rockville, MD 20852



SEP 28 1993



Sporicidin

BULLETIN NOS.
303 and 403

SPECIAL INSTRUCTIONS
FOR
CLEANING AND
DECONTAMINATION OF
SURFACES/OBJECTS
SOILED WITH
BLOOD/BODY FLUIDS
AGAINST
HIV-1 (AIDS VIRUS)

SPORICIDIN BULLETIN NO. 303:

Sporicidin Brand
Disinfectant Solution and Towlette
(EPA Reg. Nos. 8383-3 and 8383-7)

- Personal Protection:**
Persons handling items soiled with blood or body fluids likely to be contaminated with HIV-1 (AIDS virus) must protect themselves by use of disposable latex gloves, gown, mask and eye covering.
- Decontamination of Contaminated Objects:**
Immediately after use, items contaminated with blood, body fluids, or other soils that may contain HIV-1 (AIDS virus) must be thoroughly wetted with Sporcidin brand disinfectant solution or towlette for at least 1 minute before further processing.
- Cleaning Procedure:**
Following decontamination, objects and surfaces are moved to the cleaning and preparation area where they are washed; rinsed and rough dried to remove all blood and body fluids before application of the disinfectant.
- Contact Time for Disinfection:**
Thoroughly wet entire object or surface with solution or towlette. HIV-1 (AIDS virus) is killed within 1 minute contact time. *Note: Use a 10 minute contact time for other organisms. This product should not be used for disinfection of critical or semi-critical items.
- Disposal of Infectious Materials:**
Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

SPORICIDIN BULLETIN NO. 403:

Sporicidin Brand
Disinfectant Spray
(EPA Reg. No. 8383-4)

- Personal Protection:**
Persons handling items soiled with blood or body fluids likely to be contaminated with HIV-1 (AIDS virus) must protect themselves by use of disposable latex gloves, gown, mask and eye covering.
- Decontamination of Contaminated Objects:**
Immediately after use, items contaminated with blood, body fluids, or other soils that may contain HIV-1 (AIDS virus) must be sprayed to thoroughly wet with disinfectant for 1 minute before further processing.
- Cleaning Procedure:**
Following decontamination, objects and surfaces are moved to the cleaning and preparation area where they are washed; rinsed and rough dried to remove all blood and body fluids before application of the disinfectant.
- Contact Time for Disinfection:**
Spray to thoroughly wet entire object or surface. HIV-1 (AIDS virus) is killed within 1 minute contact time. *Note: Use a 10 minute contact time for other organisms. This product should not be used for disinfection of critical or semi-critical items.
- Disposal of Infectious Materials:**
Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

BEST AVAILABLE COPY