

PH 20 8383-7

10/2

FRONT PANEL

DISINFECTANT TOWELETTE

Contamination and Odor Control

SPORICIDIN  
Brand  
Disinfectant Towelette

Continuous Residual Activity  
for Over 6 Months\*

ACTIVE INGREDIENTS:

Phenol . . . . . 1.56%  
Sodium Phenate . . . . . 0.06%

- BACTERICIDAL
- FUNGICIDAL
- VIRUCIDAL\*\*
- TUBERCULOCIDAL
- ELIMINATES ODORS\*
- PREVENTS MOLD AND MILDEW

INERT INGREDIENTS: 98.38%

Solution is alkaline, pH 7.9-8.6

Keep Out of Reach of Children


CAUTION:

Avoid direct food contact. On eye contact, flush with water.  
If necessary, contact physician.

Contents:

(1) Towelette saturated  
with disinfectant.

EPA Reg. No. 8383-7



2072

BACK PANEL

DISINFECTANT TOWELETTE

**HOSPITAL DISINFECTION**

An effective disinfectant and active deodorant for hospitals, clinics, medical and veterinary offices, laboratories, industrial clean rooms, homes, nursing homes, ambulances, hotels, restaurants, schools, airplanes, trains, boats, autos, buses, health spas and toilets.

**DISINFECTANT.** Acts effectively on germ-laden non-porous equipment and surfaces in medical treatment and patient rooms, operating rooms, emergency rooms, recovery rooms and bathrooms. For use on medical equipment \*\*\* and beds, surgical carts, counter tops, manikins, telephones, furniture, wheelchairs, walkers, sinks, floors, walls, light switches, linen hampers, shower stalls, toilet seats, trash containers, damp and musty areas, including animal enclosures. Reduces the hazards of cross-contamination on environmental surfaces.

**DEODORANT.** \*It provides continuous residual bacteriostatic activity against odor-causing organisms, for over 6 months in the presence of adequate moisture if not removed from surfaces including vinyl, glass and wood.

◆Non-staining. ◆Contains no harmful abrasives.

**SPORICIDIN Brand DISINFECTANT TOWELETTES** are germicidal when used as directed for disinfection but differ in composition from other Sporcidin products.

**PASSES THE AOAC EFFICACY STANDARDS FOR INSTITUTIONAL AND HOSPITAL DISINFECTION** such as Staphylococcus aureus, Streptococcus pyogenes, Streptococcus salivarius, Salmonella choleraesuis, Pseudomonas aeruginosa, Escherichia coli, \*\*Herpes simplex types 1 and 2 (oral and genital), Influenza A<sub>2</sub> (Japan 305/57 Asian strain), and Polio type 1 viruses. HIV-1 (AIDS virus) on environmental surfaces, Trichophyton mentagrophytes (Athlete's Foot fungus) and Mycobacterium tuberculosis on environmental surfaces.

**DIRECTIONS:**

It is a violation of U.S. Federal Law to use this product in a manner inconsistent with its labeling.

**PRECLEAN** surface to remove soil and filth. Wipe dry.

**TO DISINFECT AND DEODORIZE:** Thoroughly wet surface with towelette. Ensure surface remains moist for a 10 minute contact time at room temperature (20°C/68°F or above).

**CLEAN AIR DUCTS:** Manually apply to hard non-porous precleaned duct surfaces. Rinse surfaces after application.

**TO PREVENT MOLD AND MILDEW:** Repeat application weekly, or sooner if new growth appears.

**NOTE:** See Sporcidin Bulletin No. 303 for HIV-1 (AIDS virus) instruction.

**STORAGE AND DISPOSAL;** Store at room temperature. Discard used towelettes and empty container in regular trash.

\*\*\* Other than critical and semi-critical items.

SPORICIDIN INTERNATIONAL

ACCEPTED  
17 908 1-4  
2383-7

5901 Montrose Road  
Rockville, MD 20852