



8399-6

TELONE II

Clean, Clear, Non-Clogging
SOIL FUMIGANT

for Preplant Treatment of Crop Lands to Control Plant Parasitic Nematodes and Certain Other Soil Pests

GENERAL INFORMATION

Use TELONE II soil fumigant only as a preplanting soil treatment to control plant parasitic nematodes (root-knot, meadow lesion), citrus cyst formers (golden, sugar beet, soybean), burrowing, ring, spiral, sang, pin, stubby root, stylet, dagger; also to control wireworm and garden centipedes (symphylans). Fumigate land to be planted to the crops listed below by applying TELONE II under the conditions, and at the rates recommended under DIRECTIONS FOR USE, DOSAGE AND USE RECOMMENDATIONS, and USE PRECAUTIONS. Read the entire label before using TELONE II.

- Vegetable Crops:**
- | | | | | |
|------------------|-------------|----------------|----------|-----------------|
| asparagus | cauliflower | horseradish | parsnips | shallots |
| beans | celery | kale | peas | spinach |
| beets | collards | kohlrabi | peppers | squash (summer) |
| blackeyed peas | compeas | leeks | pimentos | squash (winter) |
| broccoli | corn | lettuce | potatoes | sweet potatoes |
| brussels sprouts | cucumbers | melons | pumpkins | swiss chard |
| cabbage | egg plant | mustard greens | radishes | tomatoes |
| cantaloupe | endive | okra | rutabaga | turnips |
| carrots | garlic | onions | salsify | watermelons |
- Field Crops:**
- | | | | |
|-------------------|-----------|---------------|-------------|
| alfalfa | flax | oats | sorghum |
| barley | grasses | pasture grass | soybeans |
| birdsfoot trefoil | hops | peanuts | sugar beets |
| blackoat | lespedeza | popcorn | sugar cane |
| cover | millet | rice | tobacco |
| corn | milf | rye | vetch |
| cotton | mint | safflower | wheat |
- Citrus Fruit Tree Planting Sites:**
- | | | | |
|------------|--------|------------|----------|
| grapefruit | lemons | oranges | tangelos |
| kumquats | limes | tangerines | |
- Deciduous Fruit and Nut Tree Planting Sites:**
- | | | | |
|-------------|--------------|------------|--------------|
| almonds | dates | olives | plums |
| apples | figs | peaches | pomegranates |
| apricots | filberts | pears | prunes |
| cashew nuts | hazelnuts | pecans | quince |
| cherries | hickory nuts | persimmons | walnuts |
| chestnuts | nectarines | pineapple | |
- Bush and Vine Planting Sites:**
- | | | | |
|---------------|--------------|---------------|--------------|
| blackberries | currants | huckleberries | youngberries |
| blueberries | dewberries | loganberries | |
| boysenberries | gooseberries | raspberries | |
| cranberries | grapes | strawberries | |

Nursery Crops including floral plants, ornamentals, shrubs and bushes; forest, shade, fruit and nut trees and vine and bramble fruits of all types.

DIRECTIONS FOR USE

WHEN TO TREAT: Apply TELONE II either in the spring or fall, whenever soil type and conditions permit. For best results with annual crops, treat the soil each year. In northern states, late summer or early fall treatment (before October 15) is best for land to be planted to early spring crops, especially transplanted crops such as celery, tomatoes, and nursery and orchard stock. Early fall treatment permits planting a fall cover crop. Note: Treat muck soils only in the early fall and plant as late as possible in the spring; treat fine textured (clay) soils only when they are near or at the wilting point. Do not use TELONE II to treat any type of soil when it is cold and/or wet.

SOIL PREPARATION: TELONE II gives best results when conditions permit rapid diffusion of the fumigant through the soil and the soil surface can be sealed to prevent excessive fumigant loss during the exposure period. The soil should be in good seed bed condition, free of clods and underdecomposed plant material, moisture at about one-half of field capacity, and temperature between 40° and 80° F at the depth of injection. If underdecomposed plant debris is

ACTIVE INGREDIENT:
1,3-Dichloropropene 92%
INERT INGREDIENTS: 8%

AERATION AND PREPARATION OF SOIL BEFORE PLANTING: At the end of the exposure period allow the soil to aerate completely before planting the crop. Aeration is usually complete when the odor of TELONE II is no longer evident. Under optimum soil and weather conditions, allow one week aeration time for each 10 gallons of TELONE II applied per acre. When TELONE II is used for treating deep-rooted tree and shrub planting sites, a 3 to 6 months aeration period should be allowed. To hasten aeration, especially if heavy rains or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife-like chisel in the bed without turning the soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

DOSAGE AND USE RECOMMENDATIONS to Control Nematodes, Symphylans† and Wireworms‡

Crops (consult list of individual crops under General Information)	Type of Treatment	Soil Type	Dosage	
			Gallons Per Acre ¹	Fl. Oz./1000 ft. Row Per Chisel
Shallow Rooted Plants: Field Crops Floral Crops Grasses and Turf Small Fruits Vegetables Ornamentals	Row (42") ⁴	Mineral	4.5 to 6	46 to 62
		Muck or Peat	9 to 12	93 to 123
Strawberries	Overall (or Broadcast)	Mineral	9 to 15 ²	26 to 44
		Muck or Peat	24 ³ to 36	71 to 106
Sugar Beets Root Knot Nematode	Row (42")	Mineral	6	62
		Muck or Peat	12 to 15	35 to 44
Sugar Beet Nematode	Row (42")	Mineral	9	93
		Muck or Peat	12 to 18	35 to 53
Pineapple ⁴	Row	Mineral	24 to 36	—
Citrus — Florida ⁵	Overall	Mineral	36	106

Overall: Gallons Per Acre to Penetrate Various Depths

	3 ft.	4 ft.	5 ft.	6 ft.	
Nursery and Field: ⁷ Citrus Fruit Trees Deciduous Fruit Trees ⁸ Forest Trees Grapes Nut Trees Ornamentals (deep rooted)	Mineral Soils	3 ft.	4 ft.	5 ft.	6 ft.
	Sand	15	21	27	33
	Sandy Loam	27	30	36	48
	Silt Loam	42	51	63	75
	Clay Loam	54	69	84	102

¹ Use the higher rates in heavier soil.

Mint in Northwestern States: Use 59 gallons of TELONE II (per chisel) as an overall treatment in the spring. Reduction of the damaging effects of Verticillium wilt on mint production. After treatment allow at least 7 days to have left the soil before planting. Consult State Agricultural Extension Service Specialists for the use of other plant control, and cultural practices when using TELONE II. Verticillium wilt.

USE PRECAUTIONS

Important — Note carefully. Fumigation may form toxic and soluble salts in the soil. This is most likely when fumigant are applied to soils that are either cold or wet. Avoid injury to plant roots, fertilize as indicated to avoid ammonia injury or nitrate starvation, or both. Fertilizers containing nitrates until after the crop is above 65° F.

Certain crops including cotton, sugarcane and pomegranates above rule does not apply to them. When using in main state nursery regulations, timing of night soil application and reduce the possibility of ammonia toxicity. Citrus seedlings, Cornus sp., Crataegus sp., species have shown evidence of phosphorus deficiency to effect, it is suggested that additional phosphate be applied if a deficiency may occur.

Attention: To avoid reinfestation of treated soil, tools, seed pieces, or crop remains that could carry nematodes, etc., should be cleaned carefully before using.


Since TELONE II soil fumigant is corrosive under certain conditions TELONE II may be severe to certain materials such as rubber gloves and boots. Polyethylene provides a good barrier. Store TELONE II in tightly sealed dwellings. In outside storage, store drums on the top or bottom recessed areas.

Do not allow contamination of seeds, plants, fertilizer, feed, or domestic water supply. To avoid injury to fish and other wildlife, do not spray or other bodies of water.

Rinse equipment and containers and dispose of them from water supplies. Containers should be disposed of with wastes.

Use this product only as specified on this label.

NOTICE: Seller warrants that the product conforms to the purposes stated on the label when used under normal conditions for use. BUT NEITHER THIS WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXTENDS TO THE USE OF THIS PRODUCT OR UNDER ABNORMAL CONDITIONS OR UNDER ANY OTHER CIRCUMSTANCES. BUYER ASSUMES ALL RISK.



beets	collards	kohlrabi	peppers	squash (summer)
blackeyed peas	corn	leeks	pimentos	squash (winter)
broccoli	cowpeas	lettuce	potatoes	sweet potatoes
brussels sprouts	cucumbers	melons	pumpkins	swiss chard
cabbage	egg plant	mustard greens	radishes	tomatoes
cantaloupe	onion	okra	rutabaga	turnips
carrots	garlic	onions	salsify	watermelons

Field Crops:			
alfalfa	flax	oats	sorghum
barley	grasses	pasture grass	soybeans
birdsfoot trefoil	hops	peanuts	sugar beets
buckwheat	lespedeza	popcorn	sugar cane
clover	millet	rice	tobacco
corn	milo	rye	vetch
cotton	mint	safflower	wheat

Citrus Fruit Tree Planting Sites:			
grapefruit	lemons	oranges	tangelos
kumquats	limes	tangerines	

Deciduous Fruit and Nut Tree Planting Sites:			
almonds	dates	olives	plums
apples	figs	peaches	pomegranates
apricots	filberts	pears	prunes
cashew nuts	hazelnuts	pecans	quince
cherries	hickory nuts	persimmons	walnuts
chestnuts	nectarines	pineapple	

Bush and Vine Planting Sites:			
blackberries	currants	huckleberries	youngberries
blueberries	dewberries	loganberries	
boysenberries	gooseberries	raspberries	
cranberries	grapes	str. berries	

Nursery Crops including floral plants, ornamentals, shrubs and bushes; forest, shade, fruit and nut trees and vine and bramble fruits of all types.

DIRECTIONS FOR USE

WHEN TO TREAT: Apply TELONE II either in the spring or fall, whenever soil type and conditions permit. For best results with annual crops, treat the soil each year. In northern states, late summer or early fall treatment (before October 15) is best for land to be planted to early spring crops, especially transplanted crops such as celery, tomatoes, and nursery and orchard stock. Early fall treatment permits planting a fall cover crop. Note: Treat muck soils only in the early fall and plant as late as possible in the spring; treat fine textured (clay) soils only when they are near or at the wilting point. Do not use TELONE II to treat any type of soil when it is cold and/or wet.

SOIL PREPARATION: TELONE II gives best results when conditions permit rapid diffusion of the fumigant through the soil and the soil surface can be sealed to prevent excessive fumigant loss during the exposure period. The soil should be in good seed bed condition, free of clods and undecomposed plant material, moisture at about one-half of field capacity, and temperature between 40° and 80° F at the depth of injection. If undercomposed plant debris is present, it should be plowed down and allowed to decompose before applying TELONE II. Tillage deeper than 12 inches is necessary for good fumigant penetration in soils where a hard or plow pan occurs at plow depth. Where deep tillage is used, the tillage equipment may often be modified to simultaneously apply the fumigant and thus avoid going over the field twice.

APPLICATION: TELONE II may be applied either as an overall (broadcast) or row treatment, using suitable application equipment that will ensure placement of the fumigant at least 6 to 8 inches below the final soil surface. For overall application use either plowsole or chisel equipment with the chisels spaced 12 inches apart. When the fumigant is injected at a depth of 12 inches or more (deep tillage), the chisel spacing may be up to twice the application depth but should not exceed 30 inches. Application may be made in the same direction or at an angle to the direction of the planting row, whichever is most convenient. For row application, use chisel equipment with one chisel per row or two chisels spaced 12 inches apart to treat only the soil where the crop is to be planted. When one chisel per row is used, adjust the fumigant flow rate to distribute about 1 1/2 times more fumigant per chisel than is recommended for overall application. When two chisels are used per row, apply at the same flow rate per chisel as for overall. In both cases, the amount of fumigant required per acre will decrease as the distance between rows is increased and vice versa. At time of planting, avoid placing the seed row directly over the furrow left by the applicator chisel. When a single chisel is used per row, place the seed 3 to 4 inches to one side of the chisel furrow; when two chisels are used, plant in the center of the area between the chisel furrows.

SEALING: Immediately after application, compact the soil surface to prevent excessive fumigant loss. After chisel application, use a roller, cultipacker, or similar sealing device. After plow-sole application, disk the land, then compact it by floating or rolling. Sealing after row application can be accomplished by the tractor wheel, by listing, or by bedding so that the fumigant will be 12 to 14 inches below the top of the bed. When fumigating listed rows, seal in the fumigant with ring rollers, press sealers, or by re-listing.

EXPOSURE PERIOD: After application and sealing, leave the soil undisturbed for 7 to 14 days. A longer exposure period will be required if the soil becomes excessively cold or wet during the exposure period.

When TELONE II is used for treating deep-rooted tree and shrub planting sites, a 3 to 6 months aeration period should be allowed. To hasten aeration, especially if heavy rains or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife-like chisel in the bed without turning the soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

**DOSAGE AND USE RECOMMENDATIONS
to Control Nematodes, Symphylans† and Wireworms‡**

Crops (consult list of Individual crops under General Information)	Type of Treatment	Soil Type	Dosage	
			Gallons Per Acre ¹	Fl. Oz./1000 ft. Row Per Chisel
Shallow Rooted Plants: Field Crops Floral Crops Grasses and Turf Small Fruits Vegetables Ornamentals	Row (42") ²	Mineral	4.5 to 6	46 to 62
		Muck or Peat	9 to 12	93 to 123
Strawberries	Overall (or Broadcast)	Mineral	9 to 15 ²	26 to 44
		Muck or Peat	24 ³ to 36	71 to 106
Sugar Beets Root Knot Nematode	Row (42")	Mineral	24 to 36	71 to 106
		Overall	6	62
Sugar Beet Nematode	Row (42")	Mineral	12 to 15	35 to 44
		Overall	9	93
Pineapple ⁴	Row	Mineral	12 to 18	35 to 53
		Overall	24 to 36	—
Citrus — Florida ⁵	Overall	Mineral	36	106

Nursery and Field: ⁷	Overall: Gallons Per Acre to Penetrate Various Depths				
	Mineral Soils	3 ft.	4 ft.	5 ft.	6 ft.
Citrus Fruit Trees	Sand	15	21	27	33
Deciduous Fruit Trees ⁸		27	30	36	48
Forest Trees	Silt Loam	42	51	63	75
Grapes		54	69	84	102
Nut Trees	Clay Loam	42	51	63	75
Ornamentals (deep rooted)		54	69	84	102

- 1 Use the higher rates in heavier soil.
- 2 For cyst-forming nematodes increase dosage to 18 gallons per acre (53 fl. oz. / 1000 ft. row per chisel).
- 3 For muck soils containing less than 30% organic matter use 18 gallons per acre.
- 4 For Hawaiian pineapple, application may be made at time of or just before planting.
- 5 For burrowing nematode in citrus inject on 18-inch centers, 12 inches deep. Keep free of plants susceptible to burrowing nematodes for 2 years before replanting to citrus.
- 6 Row treatment is not recommended for potatoes in irrigated areas of west and northwestern states.
- 7 Tree planting sites prepared by backhoeing may be treated by pouring 27 fl. oz. of TELONE II into the hole during backfilling. For best results prepare and treat sites in the fall and plant in the spring.
- 8 Use TELONE II as an aid in the control of bacterial canker and decline of peach trees by application as a preplant, overall treatment of light (sandy) soils at the rate of 24 to 36 gallons per acre (71 to 106) fl. oz./1,000 ft. row per chisel) preferably in the fall when the soil is warm (50-85° F at 6-inch depth) and moist. Inject the fumigant at a depth of 10 to 12 inches with chisels mounted on 12 inch centers. Follow direction for soil preparation, sealing, exposure, and aeration as specified elsewhere on this label.

†NOTE: To control symphylans (garden centipedes) use only overall at 18 or more gallons per acre, and apply during late summer or early fall when the soil is warm. To control wireworms use the higher dosages recommended for nematodes in overall or broadcast treatments.

White Potatoes in Northwestern States: Use TELONE II as a spring or preferably a fall treatment to control quackgrass and for suppression of the damaging effects of Verticillium wilt in fields to be planted to white potatoes. Apply as an overall treatment according to the following tabular directions:

Time of treatment	Gallons per acre	Fl. Oz./1000 Ft. Row Per Chisel
Spring	17 to 25	50 to 73
Fall	25 to 34	73 to 100

GALLONS
DISTRIBUTED BY
THE GREAT WESTERN SUGAR COMPANY
P. O. Box 5308 • Denver, Colorado 80217

Fumigant are applied to soils that are either avoid ammonia injury to plant roots, fertilize as indicated above 65° F.

Certain crops including cotton, sugarcane and above rule does not apply to them. When using certain state nursery regulations, timing of night nitrification and reduce the possibility of an citrus seedlings, *Cornus* sp., *Crataegus* sp., have shown evidence of phosphorus deficiency effect, it is suggested that additional phosphorus indicates a deficiency may occur.

Attention: To avoid reinfestation of treated tools, seed pieces, or crop remains that could equipment carefully before using.


Since TELONE II soil fumigant is corrosive or fuel oil or kerosene immediately after use pumps, or other transfer equipment made under certain conditions TELONE II may be a protective clothing such as rubber gloves and material. Polyethylene provides a good barrier with polyethylene bags. Store TELONE II in 5-g dwellings. In outside storage, store drums on the top or bottom recessed areas.

Do not allow contamination of seeds, plants, contaminant food, feed stuffs, or domestic waste. To avoid injury to fish and other wildlife, do not or other bodies of water.

Rinse equipment and containers and dispose from water supplies. Containers should be disposed with wastes.

Use this product only as specified on this label.

NOTICE: Seller warrants that the product conformably fit for the purposes stated on the label under normal conditions for use, BUT NEITHER THE OF MERCHANTABILITY OR FITNESS FOR APPLIED, EXTENDS TO THE USE OF THIS PRODUCT OR UNDER ABNORMAL CONDITIONS, OR UNSEEABLE TO SELLER, AND BUYER ASSUMES



**KEEP OUT OF REACH
HAZARDOUS LIQUID
MAY BE FATAL IF INHALED,
OR SWALLOWED
OR SWALLOWED
CAUSES BURNS
OR COMBUSTION**

Do Not Get in Eyes, on Skin, or on Clothing. Necessary to Prevent Contact When Handling and Before Eating and Drinking and Air Contaminated Shoes. The POUR, SPILL, OR STORE NEAR HEAD. Adequate Ventilation. Do Not Breathe

IN CASE OF CONTACT immediately remove clothing with soap and plenty of water. FOR EYES: flush get medical attention.

IN CASE OF SPILLAGE IN DOORS have available fresh block disaster meeting specifications of a full face self-contained breathing apparatus. IF INHALED: remove to fresh air, use artificial respiration if necessary. IF SWALLOWED: call a doctor, induce vomiting or table salt in a glass of warm water.

EPA Registration No. 3399-6



8399-6

ACCEPTED

TELONE II

Clean, Clear, Non-Clogging Liquid

SOIL FUMIGANT

JUL 12 1975
FEDERAL AGRICULTURAL MECHANICAL AND CHEMICAL ACT
REGISTERED SUBJECT

Preplant Treatment of Crop Lands to Control Plant Parasitic Nematodes and Certain Other Soil Pests

ACTIVE INGREDIENT:
1,3-Dichloropropene 92%
INERT INGREDIENTS: 8%

AERATION AND PREPARATION OF SOIL BEFORE PLANTING: At the end of the exposure period allow the soil to aerate completely before planting the crop. Aeration is usually complete when the odor of TELONE II is no longer evident. Under optimum soil and weather conditions, allow one week aeration time for each 10 gallons of TELONE II applied per acre. When TELONE II is used for treating deep-rooted tree and shrub planting sites, a 3 to 6 months aeration period should be allowed. To hasten aeration, especially if heavy rains or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife-like chisel in the bed without turning the soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

DOSAGE AND USE RECOMMENDATIONS to Control Nematodes, Symphylans† and Wireworms‡

Crops (consult list of individual crops under General Information)	Type of Treatment	Soil Type	Dosage		
			Gallons Per Acre ¹	Fl. Oz./1000 ft. Row Per Chisel	
Shallow Rooted Plants: Field Crops Floral Crops Grasses and Turf Small Fruits Vegetables Ornamentals	Row (42") ¹	Mineral	4.5 to 6	46 to 62	
		Muck or Peat	9 to 12	93 to 123	
	Overall (or Broadcast)	Mineral	9 to 15 ²	26 to 44	
		Muck or Peat	24 ³ to 36	71 to 106	
Strawberries	Overall	Mineral	24 to 36	71 to 106	
Sugar Beets Root Knot Nematode	Row (42")		6	62	
	Overall	Mineral	12 to 15	35 to 44	
Sugar Beet Nematode	Row (42")		9	93	
	Overall		12 to 18	35 to 53	
Pineapple*	Row	Mineral	24 to 36	—	
Citrus — Florida*	Overall	Mineral	36	106	
Nursery and Field: ² Citrus Fruit Trees Deciduous Fruit Trees* Forest Trees Grapes Nut Trees Ornamentals (deep rooted)	Overall: Gallons Per Acre to Penetrate Various Depths				
	Mineral Soils	3 ft.	4 ft.	5 ft.	6 ft.
	Sand	15	21	27	33
	Sandy Loam	27	30	36	48
	Silt Loam	42	51	63	75
Clay Loam	54	69	84	102	

¹ Use the higher rates in heavier soil.
² For cyst-forming nematodes increase dosage to 18 gallons per acre (53 fl. oz./1000 ft. row per chisel).
³ For muck soils containing less than 70% organic matter use 18 gallons per acre.

Mini in Northwestern States: Use 50 gallons of TELONE II per acre (173 fl. oz./1000 ft. row per chisel) as an overall treatment in the spring or preferably in the fall to aid in the reduction of the damaging effects of Verticillium wilt disease. Infested and to be used to permit production. After treatment allow at least 7 to 8 weeks or until the odor of the fumigant has left the soil before planting. Consult State Agricultural Experiment Station or Extension Service Specialists for the use of other practices such as farming the land, weed control, and cultural practices when using TELONE II as an aid to reducing damage caused by Verticillium wilt.

USE PRECAUTIONS

Important — Note carefully. Fumigation may temporarily raise the level of ammonia, nitrogen and soluble salts in the soil. This is most likely to occur when heavy rates of fertilizer and fumigant are applied to soils that are either cold, wet, acid, or high in organic matter. To avoid injury to plant roots, fertilize as indicated by soil tests made after fumigation. To avoid ammonia injury or nitrate starvation, or both to crops on high organic soils, do not use fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65° F.

Certain crops including cotton, sugarcane, and pineapple are tolerant to ammonia and the above rule does not apply to them. When using high rates of TELONE II as required by certain state nursery regulations, liming of highly acid soils before fumigation may stimulate nitrification and reduce the possibility of ammonia toxicity. Certain nursery crops such as citrus seedlings, *Cornus* sp., *Gratiolus* sp., spruce and vegetable crops such as cauliflower have shown evidence of phosphorus deficiency following fumigation. To avoid this possible effect, it is suggested that additional phosphate fertilizer be used on soils where experience indicates a deficiency may occur.

Attention: To avoid reinfestation of treated soil do not use irrigation water, transplants, tools, seed pieces, or crop remains that could carry soilborne pests from infested land. Clean equipment carefully before using.

Since TELONE II soil fumigant is corrosive under certain conditions, flush all equipment with fuel oil or kerosene immediately after use. **DO NOT USE WATER.** Do not use containers, pumps, or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions TELONE II may be severely corrosive to such metals. Common protective clothing such as rubber gloves and boots, etc. may be deteriorated rapidly by this material. Polyethylene provides a good barrier. For field operations cover shoes and hands with polyethylene bags. Store TELONE II in tightly closed containers in a cool place away from dwellings. In outside storage, store drums on their sides to avoid accumulation of rain water in the top or bottom recessed areas.

Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals. Do not contaminate food, feedstuffs, or domestic water supplies.

To avoid injury to fish and other wildlife, do not spill or empty TELONE II into streams, ponds, or other bodies of water.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Use this product only as specified on this label.

NOTICE: See or warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions for use, BUT NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXPRESS OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS OR UNDER ABNORMAL CONDITIONS, OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE TO SELLER, AND BUYER ASSUMES THE RISK OF ANY SUCH USE.



KEEP OUT OF REACH OF CHILDREN

Cameraman's Note

Poor Copy

ent to control plant parasitic
olden, sugar beet, soybean),
also to control wireworm and
to the crops listed below by
amended under DIRECTIONS
E PRECAUTIONS. Read the

- shallots
- spinach
- squash (summer)
- squash (winter)
- sweet potatoes
- swiss chard
- tomatoes
- turnips
- watermelons

- sorghum
- soybeans
- sugar beets
- sugar cane
- tobacco
- vetch
- wheat

tangelos

- plums
- pomegranates
- prunes
- quince
- walnuts

youngberries

bushes; forest, shade, fruit

whenever soil type and con-
tact year. In northern states,
st for land to be planted to
r, tomatoes, and nursery and
crop. Note: Treat muck soils
ng; treat fine textured (clay)
ELONE II to treat any type of

ions permit rapid diffusion of
to prevent excessive fumigant
bed condition, free of clods
of field capacity, and tem-
dercomposed plant debris be
before applying TELONE II.
penetrative to soils where a

beans (summer)
beans (winter)
cucumbers
eggplants
melons
peas
potatoes
pumpkins
squash
tomatoes
watermelons

6 months aeration period should be allowed. To hasten aeration, especially in heavy soils or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife-like chisel in the bed without turning the soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

**DOSAGE AND USE RECOMMENDATIONS
to Control Nematodes, Symphytans† and Wireworms‡**

Crop, Common Name Individual Crop (General Information)	Type of Treatment	Soil Type	Dosage		
			Gallons Per Acre ¹	Fl. Oz./1000 Ft. Row Per Chisel	
Shallow Rooted Plants Field Crops Floral Crops Grasses and Turf Small Fruits Vegetables Ornamentals	Row (42") ²	Mineral	4.5 to 6	46 to 62	
	Overall (or Broadcast)	Muck or Peat	9 to 12	93 to 123	
Strawberries	Overall	Mineral	9 to 15 ²	26 to 44	
	Overall	Muck or Peat	24 ³ to 36	71 to 106	
Sugar Beets Root Knot Nematode	Row (42")	Mineral	6	62	
	Overall		12 to 15	35 to 44	
Sugar Beet Nematode	Row (42")	Mineral	9	93	
	Overall		12 to 18	35 to 53	
Pineapple ⁴	Row	Mineral	24 to 36	—	
Citrus — Florida ⁵	Overall	Mineral	36	106	
Nursery and Field: ⁷ Citrus Fruit Trees Deciduous Fruit Trees ⁸ Forest Trees Grapes Nut Trees Ornamentals (deep rooted)	Overall: Gallons Per Acre to Penetrate Various Depths				
	Mineral Soils	3 ft.	4 ft.	5 ft.	6 ft.
	Sand	15	21	27	33
	Sandy Loam	27	30	36	48
	Silt Loam	42	51	63	75
	Clay Loam	54	69	84	102

- Use the higher rates in heavier soil.
 - For cyst-forming nematodes increase dosage to 18 gallons per acre (53 fl. oz. / 1000 ft. row per chisel).
 - For muck soils containing less than 30% organic matter use 18 gallons per acre.
 - For Hawaiian pineapple, application may be made at time of or just before planting.
 - For burrowing nematode in citrus inject on 18-inch centers, 12 inches deep. Keep free of plants susceptible to burrowing nematodes for 2 years before replanting to citrus.
 - Row treatment is not recommended for potatoes in irrigated areas of west and northwestern states.
 - Tree planting sites prepared by backhoeing may be treated by pouring 27 fl. oz. of TELONE II into the hole during backfilling. For best results prepare and treat sites in the fall and plant in the spring.
 - Use TELONE II as an aid in the control of bacterial canker and decline of peach trees by application as a preplant, overall treatment of light (sandy) soils at the rate of 24 to 36 gallons per acre (71 to 106) fl. oz./1,000 ft. row per chisel) preferably in the fall when the soil is warm (50-85°F at 6-inch depth) and moist. Inject the fumigant at a depth of 10 to 12 inches with chisels mounted on 12 inch centers. Follow direction for soil preparation, sealing, exposure, and aeration as specified elsewhere on this label.
- [†]NOTE: To control symphytans (garden centipedes) use only overall at 18 or more gallons per acre, and apply during late summer or early fall when the soil is warm. To control wireworms use the higher dosages recommended for nematodes in overall or broadcast treatments.
- White Potatoes in Northwestern States:** Use TELONE II as a spring or preferably a fall treatment to control quackgrass and for suppression of the damaging effects of Verticillium wilt in fields to be planted to white potatoes. Apply as an overall treatment according to the following tabular directions:

Time of treatment	Gallons per acre	Fl. Oz./1000 Ft. Row Per Chisel
Spring	17 to 25	50 to 73
Fall	25 to 34	73 to 100

GALLONS
DISTRIBUTED BY
THE GREAT WESTERN SUGAR COMPANY
P. O. Box 5308 • Denver, Colorado 80217

avoid injury to plant roots, fertilize as indicated by soil tests made after fumigation. To avoid ammonia injury or nitrate starvation, or both, to crops on high organic soils do not use fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65°F.

Certain crops including cotton, sugarcane and pineapple are tolerant to ammonia and the above rule does not apply to them. When using high rates of TELONE II as required by certain state nursery regulations, liming of highly acid soils before fumigation may stimulate nitrification and reduce the possibility of ammonia toxicity. Certain nursery crops such as citrus seedlings, *Cornus* sp., *Crataegus* sp., spruce and vegetable crops such as cauliflower have shown evidence of phosphorus deficiency following fumigation. To avoid this possible effect, it is suggested that additional phosphate fertilizer be used on soils where experience indicates a deficiency may occur.

Attention: To avoid reinfestation of treated soil do not use irrigation water, transplants, tools, seed pieces, or crop remains that could carry soilborne pests from infested land. Clean equipment carefully before using.

Since TELONE II soil fumigant is corrosive under certain conditions, flush applicators with fuel oil or kerosene immediately after use. **DO NOT USE WATER.** Do not use containers, pumps, or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions TELONE II may be severely corrosive to such metals. Cotton and protective clothing such as rubber gloves and boots, etc. may be penetrated readily by this material. Polyethylene provides a good barrier. For field operations cover shoes and hands with polyethylene bags. Store TELONE II in tightly closed containers in a cool place away from dwellings. In outside storage, store drums on their sides to avoid accumulation of rain water in the top or bottom recessed areas.

Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals. Do not contaminate food, feedstuffs, or domestic water supplies.

To avoid injury to fish and other wildlife, do not spill or empty TELONE II into streams, ponds, or other bodies of water.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Use this product only as specified on this label.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions for use, BUT NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE EXPRESS OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS OR UNDER ABNORMAL CONDITIONS OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE TO SELLER, AND BUYER ASSUMES THE RISK OF ANY SUCH USE.



**KEEP OUT OF REACH OF CHILDREN
HAZARDOUS LIQUID AND VAPOR
MAY BE FATAL IF INHALED, ABSORBED THROUGH SKIN
OR SWALLOWED
CAUSES BURNS OF SKIN OR EYES
COMBUSTIBLE**

Do Not Get in Eyes, on Skin, or on Clothing. Wear Eye and Skin Protection Necessary to Prevent Contact When Handling This Material. Wash Thoroughly after Handling and Before Eating and Smoking. Wash Contaminated Clothing and Air Contaminated Shoes Thoroughly Before Reuse. DO NOT USE. POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Use Only with Adequate Ventilation. Do Not Breathe Vapor.

IN CASE OF CONTACT, immediately remove contaminated clothing and shoes and wash skin with soap and plenty of water. **FOR EYES,** flush with flowing water for at least 15 minutes and get medical attention.

IN CASE OF SPILLAGE Indoors have available an approved full face mask equipped with a fresh black canister meeting specifications of the U.S. Bureau of Mines for organic vapors, a full face self-contained breathing apparatus, or full face air-supplied respirator.

IF INHALED, remove to fresh air. Use artificial respiration if breathing has stopped. Get immediate medical attention. Keep patient absolutely quiet.

IF SWALLOWED, call a doctor. Induce vomiting by giving an emetic such as 2 tablespoons of table salt in a glass of warm water.

EPA Registration No. 8399-6

EPA Est. 8399-CO 01

Cameraman's Note
Poor Copy