

Staley
Specialty Feeds

ACCEPTED
APR 15 1976
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 8450-11

8450-11
4/5/76

P9172

RABON®

ORAL LARVICIDE

MOLASSES MINERAL BLOCK MEDICATED

TO PREVENT THE DEVELOPMENT OF HORN FLIES, FACE
FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANURE
OF TREATED CATTLE.

ACTIVE INGREDIENT

2-CHLORO-1-(2,4,5-TRICHLOROPHENYL)
VINYL DIMETHYL PHOSPHATE 0.463%

INERT* INGREDIENTS 99.537%

*Refers only to ingredients which are not insecticidal

GUARANTEED ANALYSIS

Calcium (Ca) 4.7% Min. 5.6% Max.
Phosphorus (P) 4.0% Min.
Salt (NaCl) 17.0% Min. 20.4% Max.
Vitamin A, USP Units/lb. 60,000 Min.
Vitamin D, USP Units/lb. 5,000 Min.
Total Sugars (as invert) 24.0% Min.

INGREDIENTS

Cane Molasses, Soybean Mill Feed,
Dicalcium Phosphate, Salt, Animal
Fat, BHA (a preservative), Vitamin A
Supplement, Vitamin D supplement,
Manganous Oxide, Iron Carbonate,
Copper Oxide, Cobalt Carbonate, Cal-
cium Iodate, Zinc Oxide.

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE OTHER PRECAUTIONS ON THE SIDE PANEL

EPA Reg. No. 8450-11
EPA Est. No. 8450-1L-1

NET WT. 40 LBS. — 5 BLOCKS WEIGH 200 LBS.

RABON IS THE REGISTERED TRADEMARK OF SHELL
CHEMICAL COMPANY. U.S. PATENT 3,102,812

MANUFACTURED UNDER PATENTS RE 25337 & 3,241,336

© 1976 **A. E. STALEY MFG. CO.**

DECATUR, ILLINOIS 62525

282

State

Special

Sweetlix RABON BLOCKS

PRECAUTIONARY STATEMENT HAZARDS TO HUMANS CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, consult a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. To be fed daily at the rate of .07 grams of larvicide in 0.5 ounce of block per 100 pounds of body weight. Example: 5 oz. (approximately 1/3 pound) of block for a 1000 pound animal daily.
2. Provide at least one block per five head of cattle. (So timid and smaller animals can get their share.)
3. Place the blocks where animals congregate (loafing, grazing, feeding and watering areas).
4. Feed blocks continuously. Add one additional block when EACH block has been ONE-HALF consumed.
5. Do not feed salt or mineral supplements of any kind containing salt.
6. For initial control of adult flies, spray your cattle to reduce the existing population. The normal daily intake of SWEETLIX RABON blocks will then prevent the development of new fly larvae in the manure of treated cattle. Because face flies migrate, work with your neighbors to more effectively control the adult face fly population. This product is not effective against existing adult flies. Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises. In order to achieve complete fly control SWEETLIX RABON blocks should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.