

8450-12



Staley

ACCEPTED



Staley

BEST DOCUMENT AVAILABLE
Sweetlix®



Molasses Block

TO PREVENT THE DEVELOPMENT OF HORN FLIES, FACE FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANURE OF TREATED CATTLE

ACTIVE INGREDIENT

2-CHLORO 1-(2,4,5-TRICHLOROPHENYL)
VINYL DIMETHYL PHOSPHATE 0.308%

INERT INGREDIENTS* 99.692%

*Refers only to inert ingredients which are not insecticidal

GUARANTEED ANALYSIS

Crude Protein	4.0% Min.
Crude Fat	0.5% Min.
Crude Fiber	12.5% Max.
Salt (NaCl)	19.5% Min. 23.4% Max.
Total Sugars (by weight)	20.0% Min.
Vitamin A, USP Units/lb	20,000 Min.
Vitamin D, USP Units/lb	5,000 Min.

INGREDIENTS

Cane Molasses, Soybean Meal Feed, Salt, Animal Fat, preserved w/ BHA (Butylated Hydroxyanisole), traces of Manganous Oxide, Iron Carbonate, Copper Oxide, Cobalt Carbonate, Calcium Iodate, Zinc Oxide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE OTHER PRECAUTIONS ON THE SIDE PANEL

EPA Reg. No. 645017
EPA Est. No. 645011

NET WT 33-1/3 LBS - 3 BLOCKS WEIGH 100 LBS.

A.E. STALEY MFG. CO.

Decatur, Illinois 62525

Sweetlix® R.O.L.® BLOCK

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. To be fed daily at the rate of 0.7 grams of Sweetlix R.O.L. block per 100 pounds of body weight. Example: 8 or (approximately) 1/2 pound of block for a 1000 pound animal daily. SWEETLIX R.O.L. may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.
2. Provide at least one block per five head of cattle (50 head and smaller animals range their share).
3. Place the block where animals congregate resting, grazing, feeding and watering areas.
4. Feed blocks continuously. Add one additional block when EACH block has been ONE HALF consumed.
5. Do not feed salt or mineral supplements of any kind containing salt.
6. For final control of adult flies, spray your cattle to reduce the existing population. The normal daily intake of SWEETLIX R.O.L. blocks will then prevent the development of new fly larvae in the manure of treated cattle. Because flies then migrate, work with your neighbors to more effectively control the adult fly population. This product is not effective against existing adult flies. Supplement fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or refuse on the premises. In order to achieve optimum fly control, SWEETLIX R.O.L. blocks should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

BEST DOCUMENT AVAILABLE

8450-12



ACCEPTED

Sweetlix® R.O.L.™ BLOCK

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by leaching of equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. To be fed daily at the rate of 0.7 grams of lincicide in 0.8 ounce of block per 100 pounds of body weight. Example: 8 oz. (approximately 1/2 pound) of block for a 1000 pound animal daily. SWEETLIX R.O.L. may be fed up to slaughter and to finishing dairy cows without withholding the milk from market, dairy, or other treatment.
2. Provide at least one block per five head of cattle. (So the time, smaller animals can get their share.)
3. Place the blocks where animals congregate (feeding, grazing, feeding and watering areas).
4. Feed blocks continuously. Add one additional block when EACH block has been ONE-HALF consumed.
5. Do not feed salt or other supplements of any kind containing salt.
6. For control of adult flies, spray your cattle to reduce the existing population. The normal daily intake of SWEETLIX R.O.L. blocks will then prevent the development of new fly larvae in the manure of treated cattle. Increased feed efficiency, work with your neighbors to more effectively control the adult fly population. This product is not effective against existing adult flies. Supplemental fly control measures may be needed in feed and water lots and buildings to control adult house flies and stable flies. (See block not only in cattle manure but in other decaying material, such as hay or straw on the premises. In order to achieve optimum fly control, SWEETLIX R.O.L. blocks should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage. Or disposal. Open completely to prevent. Dupes of in an incinerator or landfill approved for plastic containers, or bury in a safe place.



BEST DOCUMENT AVAILABLE

Sweetlix®

R.O.L.™

Molasses Block

TO PREVENT THE DEVELOPMENT OF HORN FLIES, FACE FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANURE OF TREATED CATTLE.

ACTIVE INGREDIENT

2-CHLORO-1-(2,4,5-TRICHLOROPHENYL)

VINYL DIMETHYL PHOSPHATE 0.308%

INERT INGREDIENTS* 99.692%

*Refers only to ingredients which are not insecticidal

GUARANTEED ANALYSIS

Crude Protein	4.0% Min.
Crude Fat	8.5% Min.
Crude Fiber	12.5% Max.
Salt (NaCl)	19.5% Min. 21.4% Max.
Total Sugars (in invert)	21.0% Min.
Vitamin A, USP Units/lb.	20,000 Min.
Vitamin D, USP Units/lb.	5,000 Min.

INGREDIENTS

Crude Molasses, Synthetic U.S. Feed Salt, Anhydrous Fat, prepared with BSA (Soyabated Hydroxypropyl), Iron Carbonate, Iron Carbonate, Copper Oxide, Cobalt Carbonate, Calcium Hydroxide, Zinc Oxide.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE OTHER PRECAUTIONS ON THE SIDE PANEL

EPA Reg. No. 8450-12

EPA Est. No. 8450-11-1

NET WT. 33-1/3 LBS. — 3 BLOCKS WEIGH 100 LBS.

A.E. STALEY MFG. CO.

Decatur, Illinois 62525

© 1980

BEST DOCUMENT AVAILABLE