

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited. Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill or bury in a safe place

FEEDING DIRECTIONS


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Adult cattle must consume 4.0 ounces per head per day to receive the 2 grams of phenothiazine intended. Young cattle must consume 1/2 ounce per 100 lbs. of body weight in order to receive the level of phenothiazine intended (1/4 gram of phenothiazine per 100 lbs. of body weight).

Free Choice: Staley Fly Control Mineral can be offered free choice to cattle even when their rations contain added mineral. Feeding stations should be located near the watering and loafing areas used by the animals.

When mixed with salt, mineral supplements, or mineral protein supplements providing salt, direct self-feeding continuously as the sole source of salt.

This product kills horn fly and face fly maggots in manure, but does not control adult flies.



Staley Fly and Worm Control Mineral with Phenothiazine

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed. Aids in the control of stomachworms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.), nodular worms (*Oesophagostomum* spp.), large-mouth bowelworms (*Chabertia* spp.), and hookworms (*Bunostomum* spp.), in cattle when fed in accordance with the directions for use.

PESTICIDALLY ACTIVE INGREDIENT

Phenothiazine.....	2.20%
Inert* Ingredients.....	97.80%
Total.....	100.0%

*Refers only to ingredients which are not pesticides.

GUARANTEED ANALYSIS

Phosphorus (P).....	minimum.....	4.00%
Calcium (Ca).....	minimum.....	16.00%
Calcium (Ca).....	maximum.....	19.00%
Salt (NaCl).....	minimum.....	12.00%
Salt (NaCl).....	maximum.....	14.00%
Vitamin A.....	minimum USP units/lb.....	50,000
Vitamin D ₃	minimum USP units/lb.....	12,500

INGREDIENTS

Calcium carbonated, monocalcium phosphate, dicalcium phosphate, soft rock phosphate, salt, iron carbonate, iron sulfate, iron oxide, manganous oxide, copper oxide, cobalt carbonate, magnesium oxide, zinc oxide, calcium iodate, vitamin A supplement, D-activated animal sterol, animal fat (preserved with BHT), and cane molasses.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

E.P.A. Reg. No. 8450-14
E.P.A. Est. No. 43807-IL-1

Manufactured for:
A. E. Staley Manufacturing Company
Decatur, Illinois 62525

Staley Fly and Worm Control Mineral with Phenothiazine

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed. Aids in the control of stomachworms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.), nodular worms (*Oesophagostomum* spp.), large-mouth bowelworms (*Chabertia* spp.), and hookworms (*Bunostomum* spp.), in cattle when fed in accordance with the directions for use.

PESTICIDALLY ACTIVE INGREDIENT

Phenothiazine.....	2.20%
Inert* Ingredients.....	97.80%
Total.....	100.0%

*Refers only to ingredients which are not pesticides.

GUARANTEED ANALYSIS

Phosphorus (P).....	minimum.....	4.00%
Calcium (Ca).....	minimum.....	16.00%
Calcium (Ca).....	maximum.....	19.00%
Salt (NaCl).....	minimum.....	12.00%
Salt (NaCl).....	maximum.....	14.00%
Vitamin A.....	minimum USP units/lb.....	50,000
Vitamin D ₃	minimum USP units/lb.....	12,500

INGREDIENTS

Calcium carbonated, monocalcium phosphate, dicalcium phosphate, soft rock phosphate, salt, iron carbonate, iron sulfate, iron oxide, manganous oxide, copper oxide, cobalt carbonate, magnesium oxide, zinc oxide, calcium iodate, vitamin A supplement, D-activated animal sterol, animal fat (preserved with BHT), and cane molasses.

KEEP OUT OF REACH OF CHILDREN
CAUTION

E.P.A. Reg. No. 8450-14
E.P.A. Est. No. 43807-IL-1

Manufactured for:
A. E. Staley Manufacturing Company
Decatur, Illinois 62525

Also adult flies may migrate in from untreated areas. Therefore, this feeding program must be used in conjunction with good management and sanitary practices; it may also be necessary to incorporate a spray program to eliminate adult flies.

WARNING: Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, avoid contact with skin and eyes. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

Do not administer to animals that are weak, anemic, emaciated, or show signs of severe constipation. Do not administer to pregnant animals in the last 4 weeks of pregnancy.

Do not use for lactating dairy animals. Occasionally an animal may be sensitive to phenothiazine resulting in symptoms such as temporary blindness and loss of skin or hair. If toxic symptoms develop, take animals off pesticide and avoid exposure to sunlight for several days.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALL STAR 12:24 MINERAL FEED (With Rabon® Oral Larvicide) may be fed free choice, top dress or mixed with ground feed. It should be consumed by cattle at the rate of .5 lbs. of mineral per 100 pounds of body weight per week (For a 600 lb. animal .90 lbs. per week, for a 1000 lb. animal 1.5 lbs. per week; for a 1400 lb. animal 2.1 lbs. per week).

This will supply the recommended average daily intake of 70 mg RABON® Oral Larvicide per 100 pounds of body weight. Animals should not be starved for minerals before initiation of the All Star 12:24 Mineral Feed program but omit all minerals from other feed after the program is started. Do not feed other free choice sources of mineral supplements in any form. Allow one mineral feeder for each 10-15 head and place near watering and loafing areas. Put out no more than one month's supply at a time and protect from rain.

If consumption of All Star 12:24 Mineral Feed is above 0.15 pounds per 100 lbs. of body weight per week, reduce the number of feeding locations or relocate. If consumption is below 0.15 pounds per 100 lbs. of body weight per week, increase the number of feeding locations. Start feeding All Star 12:24 Mineral Feed early in the spring before the flies begin to appear, and continue feeding throughout the summer and into the fall.

ALL STAR 12:24 MINERAL FEED (With RABON® Oral Larvicide) may be fed up to slaughter and to lactating dairy cows. When mixed with ground feed, care should be exercised to insure thorough and uniform mixing. When fed, it passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. Preventing larval development helps control horn flies and face flies on pasture and in files in and around cattle lots.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Pesticide which cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, ALL STAR 12:24 MINERAL FEED (With RABON® Oral Larvicide) should be used in conjunction with other good management and sanitation practices.