

## WARNING:

May be fatal if swallowed. Avoid breathing spray mist. Contact with skin can cause toxic symptoms. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

Toxic to Fish and Wild life. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not use this product in any manner not specified on this label. This product is highly toxic to bees exposed to direct treatment of residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not store or transport near food or feed. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Forage crops treated with this material should not be fed to dairy animals or to animals being finished for slaughter. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

### ANTIDOTE

**EXTERNAL** - In case of spillage on skin, wash promptly with soap and water.

**INTERNAL** - Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician IMMEDIATELY.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

WHEN CONTAINER IS EMPTY IMMEDIATELY WASH THOROUGHLY AND DESTROY. NEVER REUSE.

### DIRECTIONS FOR USE

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:  
**GROUND APPLICATION** - Vegetable and Field Crops - 30 to 250 gals.  
When Recommended for AIR APPLICATION - Field Crops - 2 to 10 gals.

**CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, TURNIPS: Cabbage Maggot** (Make preplanting soil applications at time of planting or transplanting only): Use 1 pint (1 lb. technical chlordane) in 100 gallons of water per acre.

**FIELD AND VEGETABLE CROPS (Cotton, Sorghum, Bean, Beet, Cantaloupe, Corn, Cucumber, Eggplant, Lettuce, Melons, Potato, Tomato, Cabbage): Pre-Plant Soil Treatment - BROADCAST - Wireworms, White Grubs, Cutworms, Seed Corn Maggots, Diabrotica Larvae** Use 2 to 4 pints (2 to 4 pounds technical chlordane) in sufficient water to give good coverage. Work into top 3-6 inches soil before planting.

**ANTS (Inside Buildings)** Use 5 tablespoonfuls per gallon water. Thoroughly spray or paint ant trails and areas around doorsills and window frames where the ants may enter. Treat the openings around water pipes, heat ducts, electrical outlets and baseboards where ants may come into rooms from wall spaces and partitions. Repeat applications as necessary.

**ROACHES (Inside Buildings)** Use 5 tablespoonfuls per gallon water. Thoroughly spray or paint infested cracks and other hiding places and adjacent exposed surfaces where roaches will crawl when they come out of hiding. Repeat application as necessary.

**ANTS (Outside Use)** Use 1 pint (1 lb. tech. chlordane) per 50 gallon water. Soak the colony. This chemical must be used with care around food crops. Do not apply after edible parts have formed.

**LAWNS: LAWN MOTHS (Sod Web-worms)** - 1/2 pint 2400 sq. ft. Water (soak) lawn thoroughly before lawn for 2 to 4 days after treatment. **NIGHT CRAWLERS** - 15 gals. water to 1000 sq. ft. of lawn area application. Do not allow children, pets or domestic animals to be on lawn until the grass is washed off the grass into water and grass is completely dry. **TERMITES** Use application (1 gal. per 100 gals. water) as follows:

Buildings having basements or crawl spaces: Dig a trench around foundation walls. In brick or hollow block or concrete, but not below, the footing. Then as the trench is dug, apply at rate of 4 gallons per 10 linear feet for each foot deep would require 12 gallons per linear feet.

The objective is to establish a treated soil barrier to entry. Other critical areas may also require treatment. Consult the State Extension Service for advise.

**CONDITIONS OF SALE:** 1. Cypress Mfg. Co. warrants to the chemical description on the label and as described herein. CMC neither makes, nor represents, nor warrants, any other warranty of MERCHANTABILITY, guaranteed or representation, except the material.

2. Critical and unforeseeable factors beyond CMC's control may exist in connection with the use of this material, but are not limited to, damage to plants, animals or crops. Such risks occur even though the uses stated hereon and even though the buyer and user acknowledge and assume all risks assumed by CMC under 1. above) resulting from handling this material.

Manufactured by  
**CYPRESS MANUFACTURING CO.**  
PORTAGEVILLE, MO. 63773

EPA REG. No. 8461-28

