



**ALL STAR 16:16
MINERAL FEED
(With Rabon® Oral Larvicide)**

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	70%
INERT INGREDIENTS**	99.30%
TOTAL	100.00%

*Each pound contains 3170 milligrams of Rabon®.
**Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS	
Phosphorus, min.	16.00%
Calcium, min.	15.00%
Calcium, max.	18.00%
Iodine, min.	0.01%
Salt	none added
Vitamin A, min.	100,000 USP Units/lb.
Vitamin D-3, min.	100,000 USP Units/lb.

FEED INGREDIENTS

Dicalcium and monocalcium phosphate, calcium carbonate, iron sulfate, iron oxide, potassium sulfate, magnesium sulfate, roughage product, zinc oxide, manganous oxide, copper oxide, ethylenediamine dihydrochloride, cobalt carbonate, sodium selenite, vitamin A supplement (stability improved), d-activated animal sterol (source of vitamin D-3), cane molasses and mineral oil.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back of bag for additional precautionary statements

Take Time



Observe Label - Directions
On Back Of Bag

RABON® is the registered trademark of Shell Oil Company.
U.S. Patent No. 3,102,842

ACCEPTED

APR 10 1981 J. C. PRODUCTS COMPANY, INC.
6915 Raywood Road
Madison, Wisconsin 53713

EPA Reg. No. 8487-2

EPA Reg. No. 8487-2

EPA Reg. No. 8487-2

TCP

Net Weight

50 Lbs.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALL STAR 20 MINERAL FEED (With RABON® Oral Larvicide) may be fed free choice, top dress or mixed with ground feed. It should be consumed by cattle at the rate of 15 lbs. of mineral per 100 pounds of body weight per week (For a 600 lb. animal .90 lb. per week; for a 1,000 lb. animal 1.5 lbs. per week; for a 1,400 lb. animal 2.1 lbs. per week).

This will supply the recommended average daily intake of 70 mg RABON® Oral Larvicide per 100 pounds of body weight. Animals should not be starved for minerals before initiation of the All Star 20 Mineral Feed program but omit all minerals from other feed after the program is started. Do not feed other free choice sources of mineral supplements in any form. Allow one mineral feeder for each 10-15 head and place near watering and loafing areas. Put out no more than one month's supply at a time and protect from rain.

If consumption of All Star 20 Mineral Feed is above 0.15 pounds per 100 lbs. of body weight per week, reduce the number of feeding locations or relocate. If consumption is below 0.15 pounds per 100 lbs. of body weight per week, increase the number of feeding locations. Start feeding All-Star 20 Mineral Feed early in the spring before the flies begin to appear, and continue feeding throughout the summer and into the fall.

ALL STAR 20 MINERAL FEED (With RABON® Oral Larvicide) may be fed up to slaughter and to lactating dairy cows. When mixed with ground feed, care should be exercised to insure thorough and uniform mixing. When fed, it passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. Preventing larval development helps control horn flies and face flies on pasture and those flies in and around cattle lots.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Pesticides which cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill, approved for pesticide containers, or bury in a safe place. Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control ALL STAR 20 MINERAL FEED (With RABON® Oral Larvicide) should be used in conjunction with other good management and sanitation practices.

8487-9