

**Back of Bag****W.F.A. Lawn Food with Weed & Insect Control****Front Panel**

W.F.A. Lawn Food with Weed &amp; Insect Control

W.F.A. Brand

Price - 4.95

Cover 5,000 sq. ft.

NET WT.

ORGANIC FERT.

LAWN FOOD WITH WEED &amp; INSECT CONTROL

PESTICIDE &amp; HERBICIDE GUARANTEE

**ACTIVE INGREDIENTS**

Technical Chlordane	1.238%
2,4-D Dichlorophenoxyacetic Acid	0.225%
2-(2-Methyl-4-Chlorophenoxy) Propionic Acid*	0.142%
Dicamba, 3,6-Dichloro-6-Anisole Acid**	0.060%

INERT INGREDIENTS (relative to Pesticide and Insect Content) 98.335%

\* Equivalent to 0.40% Octachlor-7-Pentanotetrahydroindane and 0.60% related compounds.

\*\* MCPP

\*\*\* Banvel®

WARNING: Keep out of reach of children. See back panel for additional caution statements.

Manufactured by:  
Western Farmers Association  
Seattle, Washington 98119

**W.F.A. Lawn Food with Weed & Insect Control** is designed to give your lawn an effective feeding with maximum control of such lawn weeds as: Dandelion, False Dandelion, Plantain, Common Chickweed, Mouse Ear Chickweed, Bealtail, White Clover, Japanese Clover, Speedwell (*Vernonia*), Buttercup, Lawn Daisies, and similar weeds. It will effectively control such harmful lawn pests as: Lawn moth, sod webworm, Billbugs, and White grubs.

**EFFECTIVE FOR APPLICATION**

1. Apply at rate of 10 to 15 pounds per 1000 sq. ft. of lawn area.
2. Spread two ways, being careful not to skip (Cyclone type spreaders not recommended for application.)
3. Apply on established lawn only. Do not apply within drip line of trees and shrubs.
4. Lawn should be dampened either from dew or light lawn watering before application.
5. Do not mow lawn for two days before or three days following application.
6. Normal lawn watering may be resumed after 24 hours.
7. Do not apply to lawn where clover is a desired part of turf. Possible injury may occur if used on St. Augustine grass, Bahiagrass, Bentgrass and Carpetgrass.
8. Occasional severe infestation of weed or insects may require repeat standard application spaced by one month.

**CAUTION**

Avoid inhalation of dust. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid contact with vegetables, flowers, shrubs, ornamentals or other plants that are sensitive to 2,4-D, Banvel® and MCPP. Do not store near growing plants, fertilizers, seeds, insecticides and fungicides. Clean spreader after each use. Wash thoroughly with soap and water after handling and before eating or smoking.

**ANTIDOTE**

External: In case of spillage on skin, wash promptly with soap and water.

Internal: Emetic of one tablespoon of mustard in tumbler of water. CALL A PHYSICIAN.

Eyes: Flush with copious amounts of water. CALL A PHYSICIAN.

**GUARANTEED FERTILIZER ANALYSIS**

6-2-4 .5% NPK

6% Nitrogen (N) Urea Formaldehyde

Ammonium Sulphate

Ferrous Ammonium Sulphate

2% Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) Single Superphosphate4% Soluble Potash (K<sub>2</sub>O) Muriate of Potash, Sul-Po-Mag

.5% Magnesium (Mg) Sul-Po-Mag

1% Iron(Fe) Ferrous Ammonium Sulphate

Organic base Sheep Manure Granular Gypsum

**CAUTION**

Keep children and pets off of treated areas until material is washed into soil and grass is dry. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.

**SPREADER SETTINGS**

SCOTT	LILLY	W.M.C. 243%
54	7	3

**NOTICE**

Western Farmers Assoc. makes no warranty, express or implied, concerning this material, except that it conforms to the chemical description on the label. Western Farmers Association disclaims responsibility in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer researches all risk and liability therefrom and accepts and uses this material on these conditions.