

ACCEPTED
 9-7-71
 8526-3 Comment



PROTEIN - VITAMIN - MINERAL SUPPLEMENT FOR RUMINANTS

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle; aids in prevention of stomach worms (*Haemonchus* spp., *Ostertagia* spp. and *Trichostrongylus* spp.) hookworms (*Bunostomum* spp.) and nodular worms (*Oesophagostomum* spp.) in cattle. Aids in prevention of foot rot in cattle and soft tissue lumpy jaw; nutritional source of iodine when fed according to the directions on this label.

PESTICIDALLY ACTIVE DRUG INGREDIENT:	
Phenothiazine	2.00%
INERT INGREDIENTS: 98.00%	
ACTIVE DRUG INGREDIENTS OTHER THAN PHENOTHIAZINE:	
Ethylene Diamine Dihydriodide	0.050%

GUARANTEED ANALYSIS

Min. Crude Protein	7.00%	Max. Calcium	2.00%
Min. Crude Fat	2.00%	Min. Calcium	1.00%
Max. Crude Fiber	4.00%	Min. Iodine	0.0357%
Max. Ash	26.00%	Max. Salt	21.00%
Min. Phosphorus	1.50%	Min. Salt	17.00%
Min. Vitamin A			50,000 USP Units/lb.
Min. Vitamin D-2			10,000 USP Units/lb.

INGREDIENTS

Wheat Millrun, Milo, Salt, Feeding Cane Molasses, Defluorinated Phosphate, Stabilized Vit. A Palmitate, Calcium Carbonate, Magnesium Sulphate, Copper Sulphate, Manganese Sulphate, Zinc Oxide, Ferrous Sulphate, Cobalt Carbonate, D-Activated Plant Sterol (Source of Vit. D-2).

FEEDING DIRECTIONS

Feed WFA's FLY CONTROL BLOCKS free choice to beef cattle. Feed at a rate of 3 1/2 oz. per 800 pounds of body weight daily. Do not feed any other salt or mineral mixture as this might lower consumption. Supply ample number of blocks - at least two for every 25 head of cattle.

CAUTION: KEEP OUT OF REACH OF CHILDREN

WARNING: Do not feed to lactating dairy animals. Occasional sensitivity to Phenothiazine has been reported in cattle. Phenothiazine is not effective against existing adult flies. Spray cattle at least once in early spring and feed WFA FLY CONTROL BLOCKS until frost. To reduce existing face fly population, a thorough larvicidal program should be carried out on an area basis. For best results cattle should be dewormed with a therapeutic dose twice per year, in the spring and the fall, for removal of specified parasites. Continuous low level treatment in conjunction with good herd management practices will control the specified parasites and will help prevent reinfestation.

USDA Reg. No. 8526-3

Net Weight 33 1/3 pounds

Manufactured for
 WESTERN FARMERS ASSOCIATION
 General Offices - Seattle, Washington

ACCEPTED WITH COMMENTS