

2521-11 PA122 143  
**BEST AVAILABLE COPY**

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS  
AND DOMESTIC ANIMALS:**

**DANGER**  
**EXTREMELY HAZARDOUS LIQUID AND  
VAPOR UNDER PRESSURE.**

**INHALATION MAY BE FATAL OR CAUSE  
SERIOUS ACUTE ILLNESS OR DELAYED  
LUNG OR NERVOUS SYSTEM INJURY.**

**DO NOT BREATHE VAPORS.**

**LIQUID OR EXCESSIVE VAPOR CAN CAUSE  
SERIOUS SKIN OR EYE INJURY WHICH MAY  
HAVE A DELAYED ONSET.**

**DO NOT GET LIQUID ON SKIN, IN EYES,  
OR ON CLOTHING.**

THIS PRODUCT CONTAINS CHLOROPICRIN WHICH MAY BE IRRITATING TO THE UPPER RESPIRATORY TRACT AND EVEN AT LOW LEVELS CAN CAUSE PAINFUL IRRITATION TO THE EYES, PRODUCING TEARING. IF THESE SYMPTOMS OCCUR, LEAVE THE FUMIGATION AREA IMMEDIATELY. EARLY SYMPTOMS OF CHLOROPICRIN OVEREXPOSURE ARE IRRITATION, RESPIRATORY DISTRESS AND VOMITING.

**OBSERVE THE FOLLOWING  
PRECAUTIONS:**

**GENERAL PRECAUTIONS**

1. Do not get in eyes, on skin, or on clothing.
2. Do not spill or discharge contents outside of areas confined for treatment.
3. Comply with all local regulations and ordinances.
4. It is advisable to supply your physician with information on Chloropicrin and Methyl Bromide. Literature is available from your dealer or distributor.
5. Obtain medical assistance at once in case of illness after exposure, and do not allow conditions which could accidentally cause further exposure until recovery is complete. (See Note to Physician.)

**RESPIRATORY PROTECTION:**

If the concentration of methyl bromide in the working area, as measured by a direct-reading detector device (such as a Draeger gas detector) does not exceed 5 ppm (20 mg/cu.m.), no respiratory protection is required. If this concentration is exceeded at any time, all persons in the fumigation area must wear protective goggles and a NIOSH/MSHA approved self-contained breathing apparatus (SCBA) or combination air-supplied/SCBA respirator (such as a U.S. Divers' Survivair or comparable device).

Under normal soil fumigation conditions, the concentration of methyl bromide in the working area will not generally exceed 5 ppm as a time-weighted average and no respiratory protection is required. However, there is the possibility of a spill or leak during soil fumigation. Therefore, a SCBA or combination air-supplied/SCBA respirator must be available and will be required for entry into an affected area in the event of a leak or spill.

**CLOTHING PRECAUTIONS:**

1. Wear loose clothing and shoes that are cleaned after each wearing. Do not wear jewelry, gloves, or tight clothing when handling. Methyl Bromide is heavier than air and may be trapped inside clothing and cause skin injury. If full-face respiratory protection is not required, wear goggles or full face shield for eye protection when handling liquid.
2. Following application, immediately remove clothing, shoes, and socks. Do not remove contaminated clothing or shoes until thoroughly cleaned and aerated. Drenched clothing cannot be adequately decontaminated.
3. Do not wear gloves of any type, or rubber protective clothing or rubber boots.
4. If liquid fumigant splashes or spills on clothing, remove them at once, as fumes will be an intolerable source of irritation.

**WARNING SIGNS: (Structural, Transportation, Space Fumigation)**

1. The applicator must placard or post all entrances to the fumigated area with signs bearing in English and Spanish: (1) the signal word DANGER/PELIGRO and the skull and crossbones symbol; (2) the statement, "Area under fumigation, DO NOT ENTER/NO ENTRAR"; (3) the date of fumigation; (4) name of fumigant used; (5) name, address, and telephone number of the applicator.
2. Do not remove a placard until the treated space is completely aerated. To determine whether aeration is complete, each fumigated area must be monitored and shown to contain less than 5 ppm methyl bromide. If less than 5 ppm methyl bromide is detected, the placard may be removed.
3. Placards must be placarded with a G.T. specified warning sign. Warning signs are available from your dealer or distributor.

**SPILL OR LEAK PROCEDURE:**

1. Evacuate the immediate area of the spill or leak. Use SCBA or combination air-supplied/SCBA respirator for entry into the affected area to correct problem. Move leaking or damaged cylinders or containers outdoors or to an isolated location. Following strict safety precautions, suck up spill if possible. Allow spill to be unattended until the concentration of methyl bromide is determined to be less than 5 ppm (20 mg/cu.m.) and chloropicrin less than 0.1 ppm.
2. Contaminated soil, water, and other cleanup debris is a toxic hazard. A spill report shall be filed to the National Hazardous Waste Reporting System if reportable quantity of 1000 lbs. is exceeded. (40 CFR 302.61) If reportable quantity of 1000 lbs. is exceeded,

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**PIC-BROM 25**

**PREPLANT  
SOIL FUMIGANT**

**ACTIVE INGREDIENTS:**

METHYL BROMIDE 75%

CHLOROPICRIN 25%

INERT INGREDIENTS: 0%

TOTAL 100%

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**



**PELIGRO**

**POISON**

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta la haya sido explicado ampliamente.

**IN ALL CASES OF OVEREXPOSURE, GET  
MEDICAL ATTENTION IMMEDIATELY  
TAKE PERSON TO A DOCTOR OR TO AN  
EMERGENCY TREATMENT FACILITY**

**STATEMENT OF PRACTICAL  
TREATMENT**

**IF INHALED:** Get exposed person to fresh air. Keep warm. Make sure person can breathe freely. If breathing has stopped, give artificial respiration. Give oxygen if needed. Do not give anything by mouth to any unconscious person. Seek medical attention.

**IF ON SKIN:** Immediately remove contaminated clothing, shoes, and other items covering the skin. Wash contaminated skin area thoroughly with soap and water.

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes.

**NOTE TO PHYSICIAN:** Early symptoms of overexposure are dizziness, headache, nausea and vomiting, weakness, and collapse. Lung edema may develop in 2 to 48 hours after exposure, accompanied by cardiac irregularities; these effects are the usual cause of death. Repeated overexposure can result in blurred vision, ataxic gait, and mental imbalance, with probable recovery after a period of no exposure. Do not give oxygen to suggest the occurrence, but not the degree of exposure. Treatment is symptomatic.

### PRECAUTIONS FOR SOIL FUMIGATION USE

- PRECAUTIONS TO FUMIGATION:**
1. Comply with all local regulations and ordinances. Obtain an application permit from Agricultural Regulatory Agencies as required.
  2. Never fumigate alone. It is imperative always to have an assistant and proper protective equipment in case of accidents.
  3. Drivers of application equipment must advise other workers of all precautions and procedures. In addition, drivers must instruct their helpers in the mechanical operation of the tractor and how to safely work with the tractor and driver while fumigating.
  4. Handle this fumigant in the open with the operator "up wind" from the container where there is good ventilation.
  5. Check fumigant pressure system for leaks before beginning operation.
  6. Two trained persons must be present during introduction of the fumigant.
  7. When fumigating soil from a tractor, 5 gallons of water must be carried on the tractor and placed where it is readily accessible. In addition to water available on the tractor, at least 5 gallons of additional water must be available from the service truck. This water must be potable and in containers marked "decontamination water not to be used for drinking." Field should be reasonably free of trash before starting the fumigation.
- SEE PRECAUTIONS CONTINUED IN THIRD COLUMN



Soil  
Chemicals  
Corporation  
PRODUCTS

P.O. BOX 782 · HOLLISTER, CA 95024

E.P.A. EST. 8536-CA-1,2,3,4:FL-1

E.P.A. REG. NO. 8536-11

NET CONTENTS . . . . . LBS.



**PRECAUTIONS, CONTINUED:**

- STOPPING FUMIGATION:**
1. This fumigant should not be applied when there is an atmospheric inversion. Since uncomfortable concentrations of ethylene dichloride may drift to nearby areas, immediately cover treated area with plastic tarpaulin for a minimum of 48 hours when the injection depth is less than 10".
  2. Do not lift in action chains to turn at the end of a pass until fumigant has drained from system following closure of shut-off valve.
  3. If trash is inadvertently pulled by the shanks to the ends of the field when flogging, it must be covered by lifting and shanks, then covering the trash with polyethylene film before making the turn for the next pass.
  4. When changing the cylinders, be certain they are turned off and fumigant system is not under pressure. Do not open the system when there are people or structures downwind where exposure above the permissible exposure levels could occur.

- FOLLOWING FUMIGATIONS:**
1. Post all treated areas with warning signs, available from your dealer or representative.
  2. No children, unauthorized people, or animals should be in the fumigation area for 48 hours.
  3. When tarpaulin is used, two trained persons must be present during removal of the tarpaulin.

- SPILL AND LEAK PROCEDURES:**
1. In case of a rupture of hose or fitting while applying fumigant, immediately stop tractor and motor. Evacuate immediate area of spill or leak. Use SCBA or similar air-supply/SCBA respirator for entry into affected area to correct the problem. Approach from upwind to make necessary repairs.
  2. Do not reenter area without respiratory protection until spill has evaporated or leak has been fixed.

**NOTE CAREFULLY:** Fumigation may temporarily raise the level of ammonia nitrogen and soluble salts in the soil. This is most likely to occur when heavy rates of fertilizer and fumigant are applied to soils that are either cold, wet, acid, or high in organic matter. To avoid injury to plant roots, fertilizer should be injected by soil tests made after fumigation. To avoid ammonia injury and/or nitrate starvation to crops, avoid using fertilizers containing ammonium salts and use only fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65 degrees F. Liming highly acid soils before fumigation stimulates nitrification and reduces the availability of ammonia toxicity.

**PRECAUTIONS FOR STRUCTURAL, TRANSPORTATION, OR SPACE FUMIGATION USE**

- GENERAL PRECAUTIONS:**
1. Keep animals, children, and unauthorized people away from area under treatment until area is certified free of methyl bromide (See Aeration Section).
  2. When used for fumigation of enclosed spaces (houses and other structures, warehouses, vaults, chambers, greenhouses, tractors, vans, boxcars, ships, and other transport vehicles, and tarpaulin-covered areas), two persons trained in the use of this product must be present during introduction of the fumigant. Initiation of aeration, and after aeration when testing is necessary. Two persons do not need to be present if monitoring is conducted routinely (outside the area being fumigated).
  3. Do not fumigate with this product when the temperature is below 40 degrees F.
  4. Whenever possible, apply methyl bromide from outside of structure or car being fumigated. Make sure the fumigated area is properly sealed and posted. Do not move trucks, trailers, or vans during fumigation. They must be completely averted before movement is allowed.

- AERATION AND REENTRY:**
1. After fumigation, treated areas must be aerated until the level of methyl bromide is below 5 ppm (20 ug/cu.m).
  2. Do not allow entry into the treated area by any person before this time unless loose clothing and a respiratory protective device (SCBA or combination air-supplied/SCBA) is worn.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in any manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE AND HANDLING:**  
Store in dry, cool, well-ventilated area under lock and key. Do not store in a pesticide storage area. Do not contaminate water, food, or feed by storage. Persons moving or handling containers should wear protective clothing. Open container only in a well-ventilated area wearing protective clothing, and respiratory protection if necessary. Store cylinders upright, secured to a rack or wall to prevent tipping. Cylinders should not be subjected to rough handling or mechanical shock such as dropping, bumping, dragging, or sliding. Do not use ropes, chains, hooks, lumps, or similar devices to unlash cylinders. Transport cylinders using hand truck, fork truck, or other device to which the cylinder can be firmly secured. Do not remove valve protection bonnet and safety cap until immediately before use. Replace safety cap and valve protection bonnet when cylinder is not in use. When cylinder is empty, close valve, screw safety cap onto valve outlet, and replace protection bonnet before returning to shop. Only the manufacturer is authorized to refill cylinders. Do not use cylinders for any other purpose. Follow registrant's instructions for return of empty or partially empty cylinder.

**RETURN OF CYLINDERS:**  
Cylinders are the property of:  
Soil Chemicals Corporation      Soil Chemicals Corporation  
8770 Hollister 25                      2066 East 220th Street  
Hollister, CA 95023                      Long Beach, CA 90810  
and should be returned promptly by collect auto freight. Do not ship cylinders without safety caps or valve protection bonnets. When a cylinder is partially full and there is no further requirement for the product, contact SCC for return instructions.

**GENERAL INSTRUCTIONS**

This fumigant is a highly hazardous material and should be used only by trained and trained in its proper use before using. Read and follow all label directions and directions including the attached sufficient. All persons working with this fumigant must be knowledgeable about the hazards and trained in the use of required respiratory equipment and detector devices, emergency procedures, and proper use of the fumigant.

**GENERAL INFORMATION:** This fumigant is a preplant soil fumigant for control of soil-borne pests and diseases caused by certain species of *Phytophthora*, *Fusarium*, and *Phytophthora*. Wireworms, nematodes (including *Meloidogyne*, *Pratylenchus*, *Tylenchus*, *Heterodera*, *Citricheles*, and *Paratylenchus* species), broadleaf and grassy weeds (including quackgrass, annual bluegrass, crownbeetle, common lambsquarters, tarweed, and tarweedgrass). This fumigant is not effective against mallow, filaree, dodder, some species of clover, and certain other hard seeded weeds. It is to be used in soil to be planted to the crops listed below.

**SOIL PREPARATION:** Prior to application, the soil should be in condition for planting and with sufficient moisture to support seed germination. When fumigating for the control of ratsedge, the soil should be ripped to a depth of at least 15 inches.

**APPLICATION:** Apply this fumigant by means of chisels spaced from 12" to 18" apart and 5 to 8 inches below the soil surface. To seal the fumigant after application, disc, or cultipack immediately behind chisels and then cover with a gas-tight tarpaulin.

**DOSE TABLE:**

Land To Be Planted To:	Rate, Ppm Per Acre 1,2
Broccoli	250-375
Cauliflower	250-375
Eggplants	250-375
Floral Crops	300-400
Mushrooms	250-375
Nursery Crops	300-400
Peppers	250-375
Pineapples	250-375
Plant and Seed Beds	300-825
Strawberries	250-375
Tomatoes	250-375

1. Row or bed applications may be made at the broadcast rates, but the amount used will be proportionately less per acre depending on the row spacing and width of treatment in the row or bed.
2. Use the higher rates for nutsedge control.
3. Plants grown for transplanting.

**AERATION:** Remove tarpaulin after 48 hours. Allow the soil to aerate for 2 weeks before planting. If odors of fumigant persist beyond this two-week period, disc or plow the soil to help aeration.

**DECIDUOUS FRUITS, NUTS, CITRUS, VINEYARDS AND OTHER PERENNIAL WOODY PLANTS**

**APPLICATION:** (Note: Do not harvest crop for 24 months after fumigation.) Prior to planting, apply methyl bromide at a rate and exposure time that is indicated in the dosage instructions by means of chisels spaced not more than 5 1/2 feet apart. Inject at a depth of not less than 20 inches. To seal fumigant after application for disease control, cover immediately with a gas tight tarpaulin by means of a mechanical tarp layer. The soil should be ripped to not less than 30 inches and the subsurface should be as dry as possible. Excess moisture will block deep penetration of the methyl bromide.

Clay soils and soils that drain poorly may be dried out by planting early grass and then withholding water during the summer. This will increase penetration of methyl bromide.

Back hoeing and sugaring are alternative methods of applying methyl bromide to tree sites. Attach a methyl bromide dispenser to the cylinder of methyl bromide and to it attach a polyethylene applicator tube. The correct dosage can be measured with the dispenser and then applied through the tube.

Test soils and soils very high in organic content should not be fumigated to control *Armillaria* galls with methyl bromide.

**USE:** Deciduous fruits, nuts, citrus, vineyard and other woody plants for control of *Armillaria* root rot; *Phytophthora* root rot; Crown rot fungus; Citrus nematodes; Burrowing nematode; Root Knot Nematode; Root Rot Nematode.  
**RATE:** 400 - 850 lb/acre. Minimum exposure 5 to 7 days.

**USE:** Tree Sites for control of *Phytophthora* root rot, *Armillaria* root rot in light sandy soils and fine textured clay soil.  
**RATE:** 1 to 1 1/2 lbs/100 sq. ft.

**PLANTING:** In cold climates, treat in fall and plant the following spring. In warm climates, do not plant until 30 days after application.

**FOR USE IN FLORIDA CITRUS CONTROL OF PHYTOPHTHORA IN SANDY SOILS**

This is preplant or replant treatment. Trees which are planted in this treated soil will not bear harvestable fruit for a period of at least 18 months. Use a minimum of 1 to 1 1/4 pounds per 100 square feet. Apply to fumigation for 48 hours. Covering treated area with a 4 mil tarpaulin will control disease to a depth of 4 feet. Aerate 2 weeks before setting transplants in treated area.

**FOR USE IN CALIFORNIA ONLY ASPARAGUS AND LETTUCE**

**SHIPPING:**

This fumigant is classified in the U.S. Department of Transportation Hazardous Materials Regulations as Methyl Bromide and more than 2% Chloropicrin mixture, class "B" poison (inhalation hazard) and no exceptions from specifications, packaging, marking, or labeling are allowed. Describe empty cylinders as having contained Methyl Bromide and more than 2% Chloropicrin mixture (inhalation hazard). Do not ship with foods, feeds, or clothing.

**DISPOSAL:**

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**APPLICATION:** Follow application directions as described above.

CROP	DOSAGE AND USE TABLE	
	USE	RATE
Asparagus	Fungus diseases (deeping off, Phythium spp) Weeds (grasses), Nematodes	400-530 lbs/A
Lettuce	Weeds (including scallia) Nematodes Big Vein	400-530 lbs/A 270-400 lbs/A

**APPLICATION:** Remove tarpaulin after 48 hours

**REMARKS:** For use in Calif only. Do not harvest asparagus during year of treatment. Use on lettuce fields only once in 2 to 3 years.

**ENVIRONMENTAL HAZARD**

This pesticide is toxic to wildlife. Do not discharge effluents containing this product into lakes, streams, ponds, estuaries, oceans, or public waters unless this product is specifically identified and addressed in an NPDES permit. Do not discharge effluents containing this product to sewer systems without previously notifying the sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

**CHEMICAL HAZARD**

Do not use containers or application equipment made of magnesium, aluminum, or their alloys, as under certain conditions this fumigant may be severely corrosive to such metals. Do not permit water to be used to clean fumigant pressure system as corrosion will result. Diesel oil is satisfactory for this purpose.

**WARRANTY**

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CA 2 88 - MP 10379

2063

## INSTRUCTIONS

... AND SHOULD BE ...  
BEFORE USING ...  
INCLUDING THE ATTACHED ...  
... IF RESPIRATOR IS ...  
... AND PROPER USE OF THE

A persistent soil fumigant for diseases caused by certain ... and Phytophthora. ... Pratylenchus. ... species. ... and annual bluegrass, ... sedge, and ... In ... dodder. ... species of weeds. It is to be used in ...

The soil should be in condition to support seed germination. ... the soil should be ...

... of chisels spaced from 12 to ... surface. To seal the fumigant immediately behind chisels ...

... Per Acre 1.2
250-375
250-375
250-375
250-400
250-375
250-375
250-375
250-375
250-375
250-375
250-375
250-375
250-375
250-375
250-375

... made at the broadcast ... be proportionately ... row spacing and width ... edge control

... Allow the soil to aerate for ... beyond this ...

## CITRUS, VINEYARDS WOODY PLANTS

... for 24 months after fumigation ... at a rate and exposure ... by means of chisels ... at a depth of not less ... application for disease control ... by means of a ... to not less than 30 inches ... Excess moisture will ...

... may be dried out by planting ... during the summer. This will ...

... methods of applying methyl ... dispenser to the cylinder ... polyethylene applicator tube ... the dispenser and then applied

... content should not be ... methyl bromide.

... vineyard, and other woody ... liaria root rot; Phytophthora ... Citrus nematode; Burrowing ... root knot nematodes ... exposure 5 to 7 days

Phytophthora root rot. Apply to soils and fine textured clay

... and plant the following ... 30 days after application

## FOR USE WEST OF ROCKY MOUNTAINS ONIONS, DIRECT SEEDED

APPLICATION. Treatment can be made whenever soil conditions are suitable. In northern states, late summer or early fall treatments are best for land to be planted to early spring crops. Follow application directions as described above.

USE for control of White Rot, weevils (grasses) and Nematodes

DOSE 400 lbs/Acre

RESTRICTIONS. Use restricted to west of the Rocky Mountains. Do not treat any field more than once every twelve months.

## GREENHOUSE FUMIGATION

This fumigant may be used in field or in greenhouse. When used in greenhouse, the following safety precautions must be strictly observed:

BEFORE FUMIGATION: The use of methyl bromide in confined spaces presents a potential hazard to humans and plant life. Special precautions must be made in order that these potential hazards be minimized. It is the responsibility of the individual supervising the fumigation operation to see that all safety precautions are strictly observed, before the fumigation operation commences. A supervisor of the fumigation job shall have conducted proper training of all personnel involved in the fumigation (including use of safety equipment), removed all persons from the area not directly involved in the fumigation, and inspected the equipment to insure a proper operation.

DURING FUMIGATION: If a wind is blowing, all detections should be made upwind from a previous injection site. Immediately after injection of the fumigant and tarping, a qualified person wearing protective equipment should monitor the area with a halide leak detector. If excessive leaks are found, the source of the leak should be resealed immediately. During this operation, all windows and doors should be open and fans operating to assist ventilation.

EXPANDING OF AREA: The fumigated area must be placarded (see directions under Warning Signs). Do not re-enter warning signs until the fumigated area is completely aerated and safe for entry as indicated by a suitable detector.

PROTECTIVE EQUIPMENT: Although this fumigant contains chloroform, the absence of chloroform does not always indicate the absence of methyl bromide. Under no circumstances shall any person be allowed to enter the fumigated structure without the appropriate protective equipment from the time of injection of the fumigant until acceptable readings are obtained using an approved detector. To maintain adequate safety standards, the following equipment must be present on the site during the entire fumigation operation: (1) One or more self-contained breathing apparatus; (2) One or more replacement air bottles per breathing apparatus; (3) One or more halide leak detectors.

AERATION AND REENTRY. At the end of the exposure period, remove all seals and open all doors and windows that are operational. Allow for complete ventilation. Use ventilation fans wherever possible to remove fumigant from dead air pockets. After fumigation, treated areas must be aerated until the level of methyl bromide is below 5 ppm. Do not allow entry into the treated area by any person before this time unless provided with loose clothing and a respiratory protection device (SCBA or combination air-supplied SCBA). Note: Certain materials absorb methyl bromide during fumigation, and desorption during aeration may call for extended monitoring and aeration periods.

HALIDE LAMP DETECTOR. BEHREND-HANSEN DETECTOR: (Hard Copy) and Detector Tube: Methyl Bromide may be detected at the Threshold Limit Value (TLV) of 5 ppm. Detectors are available from Soil Chlorine Corporation.

FLAME LAMP DETECTOR. Color of the flame is an index of the concentration of methyl bromide present. The following tabulated information gives an index of flame color and concentration of methyl bromide present.

MEASUREMENT (PPM)	LB/1000 CU FT	FLAME COLOR (DAYLIGHT*)
0	0	None
25	0.052	None
50	0.104	Very Light Blue
125	0.26	Green
250	0.52	Strong Green
500	1.04	Strong Blue-Green Flame
1000	2.08	Strong Blue-Green
1000	2.08	Blue

\* Nighttime color is identical but allowance must be made for the bluish cast of the flame itself.

Note: Halide lamp detector is suitable for locating leaks and for determining extent of aeration down to approximately 50 ppm. It is not suitable for clearing a structure for reentry.

IRIDA  
GROW IN SANDY SOILS

rows which are planted in this  
suit for a period of at least  
pounds per 100 square feet  
ing treated area with a 4 mil.  
th of 4 feet. Aerate 2 weeks  
a.

IRIDA ONLY  
LETTUCE

as described above

RATE  
20-510 lbs/A


0-530 lbs/A

0-400 lbs/A

not labeled  
in lettuce fields

not conform to its chemical  
e purposes stated on the label  
under normal conditions of use.  
warranty of MERCHANTABILITY or  
as or implied, extends to the  
nstructions, or under abnormal  
sonably foreseeable to seller.  
se

CA 2 88 - MP 10379



ACCEPTED  
8536-11  
20 SEP 1988

UNDER THE FEDERAL INSECT CONTROL ACT AND FEDERAL FOOD, DRUG AND COSMETIC ACT, THIS PRODUCT IS SUBJECT TO THE FOLLOWING REGULATIONS: 40 CFR 158.101-158.102. SUBJECT TO ATTACHED COMMENTS.

NA 1581  
METHYL BROMIDE  
AND MORE THAN  
2% CHLOROPICRIN  
MIXTURE, LIQUID