


# Sani-Clor Brand Algicide

BEST DOCUMENT AVAILABLE



## ACTIVE INGREDIENT:

Poly(oxyethylene (dimethyliminio) ethylene (dimethyliminio) ethylene dichloride) 60.0%

INERT INGREDIENT 40.0%

This product contains 5.76 lb. of active ingredient per gallon and weighs 9.6 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

## CAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

**FIRST AID:** If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**ENVIRONMENTAL HAZARDS.** This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency

## DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Sani-Clor Brand Algicide is used to control the growth of algae in swimming pools, exterior spas, whirlpools, hot tubs, and decorative fountains. For maximum effectiveness pools, spas, whirlpools, hot tubs, and decorative fountains containing heavy growth of algae should be cleaned prior to using Sani-Clor Brand Algicide.

For pools having just visible algae growth add an initial dose of 11 to 17 fluid ounces of Sani-Clor Brand Algicide per 10,000 gallons of water and remove settled algae debris by cleaning. For treatment of a freshly cleaned and filled pool add initially 6 to 11 fluid ounces of Sani-Clor Brand Algicide per 10,000 gallons of water. Subsequent additions of 2 to 4 fluid ounces of Sani-Clor Brand Algicide per 10,000 gallons of water should be made every 5 to 7 days after initial treatment for maintenance.

Spas, whirlpools, hot tubs, or decorative fountains having just visible algae growth require an initial dose of 1 to 2 fluid ounces of Sani-Clor Brand Algicide per 1000 gallons of water. For treatment of a freshly cleaned and filled spa, whirlpool, hot tub, or decorative fountain, add initially 0.6 to 1.1 fluid ounces of Sani-Clor Brand Algicide per 1000 gallons of water. Subsequent additions of 0.2 to 0.4 fluid ounces of Sani-Clor Brand Algicide should be made every 5 to 7 days after initial treatment for maintenance.

Sani-Clor Brand Algicide is compatible with those chemicals normally used to treat pools and spas and is effective at both acid and alkaline pH. Sani-Clor Brand Algicide can be used in pools and spas treated with chlorine chemicals and may reduce the amount of those chemicals normally required. However, do not mix Sani-Clor Brand Algicide with concentrated dry or liquid chlorine products.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Do not stack more than four drums high. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep containers closed when not in use.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL

**PLASTIC:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**METAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Manufactured by:

GPS Industries

13280 Amar Road

City of Industry, CA 91746

EPA REG. NO. 8544 6

EPA EST. NO. \_\_\_\_\_

NET CONTENTS \_\_\_\_\_