

CONTROL

HY

GIA

-3

DIRECTIONS FOR USE

DEGRADES WITH AGE. USE A TEST KIT AND INCREASE DOSAGE AS NECESSARY TO OBTAIN REQUIRED LEVEL OF AVAILABLE CHLORINE.

SWIMMING POOL CHLORINATION

For Freshly Filled Pools – Add 1 pint of product per 10,000 gallons of water. Repeat the above dosage until a chlorine residual of 0.6 to 1.0 ppm is established as determined by use of a chlorine test kit. Thereafter, add 8 ounces of product for each 10,000 gallons of water daily or as needed to maintain a chlorine residual of 0.6 to 1.0 ppm at all times. Test pool water frequently with a suitable chlorine test kit.

BATHHOUSE SANITIZATION

Diving boards, locker room benches, bath mats and shower room floors should be treated with 800 ppm (1 ounce product per gallon of water) available chlorine. Walkways in the pool area should be treated daily with 500 ppm available chlorine.

WARNING

HARMFUL IF SWALLOWED
IRRITATING TO EYES, SKIN

ANTIDOTE

EXTERNAL

WASH WITH WATER

INTERNAL

DRINK MUCILAGE, RAW EGGS, RICE GRUEL. FOLLOW WITH 1-2 SPOONFUL OF MUSTARD IN WATER

CALL A PHYSICIAN

STORAGE

DO NOT store sodium hypochlorite for more than four months. Keep container in a cool, dry place as light and heat reduce strength.

WARNING

DO NOT mix sodium hypochlorite with acids or acid detergents.

CONTAINER DISPOSAL

Return to drum reconditioner or to manufacturer for recycling or crushing and burying

DIRECTIONS FOR USE

GRADES WITH AGE. USE A TEST KIT AND INCREASE DOSAGE AS NECESSARY TO OBTAIN REQUIRED LEVEL OF AVAILABLE CHLORINE.

SWIMMING POOL CHLORINATION

Freshly Filled Pools — Add 1 pint of product per 10,000 gallons of water. Repeat the above dosage until a chlorine residual of 0.6 to 1.0 ppm is established as determined by use of a chlorine test kit. Thereafter, add 8 ounces of product for 10,000 gallons of water daily or as needed to maintain a chlorine residual of 0.6 to 1.0 ppm at all times. Test pool frequently with a suitable chlorine test kit.

BATHHOUSE SANITIZATION

Locker room benches, bath mats and shower floors should be treated with 800 ppm (1 ounce per gallon of water) available chlorine. Walkways in pool area should be treated daily with 500 ppm available chlorine.

WARNING:

HARMFUL IF SWALLOWED
IRRITATING TO EYES, SKIN AND LUNGS

ANTIDOTE

EXTERNAL

WASH WITH WATER

INTERNAL

DRINK MUCILAGE, RAW EGG WHITE, MILK OR RICE GRUEL. FOLLOW WITH EMETIC (TABLE-SPOONFUL OF MUSTARD IN GLASS OF WATER).

CALL A PHYSICIAN

STORAGE

DO NOT store sodium hypochlorite longer than four months. Keep in cool, dark place as light and heat will reduce strength.

WARNING!

DO NOT mix sodium hypochlorite with acids or acid detergents.

CONTAINER DISPOSAL

Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

REGISTERED

Jul 19 1948

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. SUBJECT
 TO ATTACHED COMMENTS.

DIRECTIONS FOR UDDER WASHING AND SANITIZING . . .

1. Prepare the udder washing solution in plastic pail. (do not use tin or galvanized pails) using 1/2 ounce Agway Iodophor to 2 1/2 U.S. gallons warm water, 110 to 115° F.
2. Approximately one minute before application of the machine, wash and massage the teats and udder with this solution, using a paper towel which has been soaked in the solution.
3. After gentle washing and massage of the udder, press the excess solution from the towel and use the reverse side to dry the teats and udder.
4. Discard the used towel, and use a clean towel for each cow.
5. Check udder with a strip cup or screening paddle before applying the machine.
6. Apply teat cups after sanitizing by dipping in a solution of 25 ppm available iodine. Surfaces and equipment to be sanitized should be exposed to the sanitizer for a period of 2 minutes.

FOR CLEANING UTENSILS AND EQUIPMENT . . .

1. Pre-rinse utensils and equipment whenever possible.
2. Wash with Agway Iodophor, using 1 ounce to each 5 gallons of warm water.
3. Rinse with clean water.
4. Sanitize in a solution of 1 oz. of Agway Iodophor, to 5 gallons of water. Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on rack or drain board to air dry. Change washing and sanitizing solutions when the brown iodine color disappears. Make fresh solution each time used.

TABLE OF DILUTIONS . . .

1 ounce Agway Iodophor to 2 1/2 U.S. gallons	50 ppm available iodine
1/2 ounce Agway Iodophor to 2 1/2 U.S. gallons	25 ppm available iodine

CAUTION . . .

Agway Iodophor is a concentrated product. Harmful if swallowed. Avoid contamination of foods. Avoid contact with skin and eyes. In case of contact, flush skin or eyes with plenty of water; for eyes get medical attention. Use in recommended dilutions.

