

ACTIVE INGREDIENTS

Captan (N-trishli 10.0%

Parathion (O,O-diethyl O-p-nitrephenyl thiophosphate) INERT INGREDIENTS.

100.0%

2.0%

38.0%



POISON W



DANGER:

KEEP OUT OF REACH OF CHILDREN.

Poisonous if swallowed, inhaled, or absorbed through skin. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for parathion protection. Rapidly absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing, and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep out of reach of children. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have discipated. Do not contaminate feed and foodstuffs. During application avoid drift to adjacent food and forage plantings which might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is highly toxic to Bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by ng of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse this empty container. Dispose of by burying with wastes or burning. Stay away from smoke and fumes.

ANT!DOTES: If swalloweds Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN WANEDIATELY! If on skin: In case of contest, ranges contaminated clathing and immediately week skin with seap and water.

Est Soiler warrants that the product conforms to its scription and is reasonably fit for the purpose stated on the label when week in accordance with directions under normal conditions of use, but neither this warranty nor any other warrenty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseemble to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include authorizential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Apples: To aid in control of scab, Brook's fruit spot, sooty blotch, fly speck, codling math, plum curculto, leaf rollers, bud moth, redbug, gahide and mites, apply at the rate of 30-50 pounds per acre using the higher rate to control bud moth and redbug. Full-coverage applications by airplane or ground duster is necessary for satisfactory control.

Repeat application at 4 to 7 day intervals as needed to maintain cover when apple scab is prevalent in the pre-bloom, bloom, petal-fall, first and second cover dust application. Continue cover sprays at 7 to 14 day intervals during growing season as necessary for disease and insect control. Do not apply during blossom period.

CAUTION: Do not apply within 14 days of harvest. Remove residues remaining on fruit at harvest by washing, wiping, or other effective means. Do not apply within 10 days of an oil spray. Use cautiously on Red Delicious and Baldwin varieties.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Agway Inc., Chemical Division Box 1333, Syracuse, N. Y. 13201

EPA Reg. No. 8590-2