



# sanitizer and teat dip

EPA Reg. No. 8590-4

**CAUTION KEEP OUT OF  
REACH OF CHILDREN**

see additional precautions  
on side panel.

### active ingredients

sodium hypochlorite . . . . . 8.4%

inert ingredients . . . . . 91.6%

**NET CONTENTS 1 U.S. GALLON**

## CAUTION

IRRITATING TO EYES,  
SKIN AND LUNGS  
HARMFUL IF SWALLOWED

**ANTIDOTE  
EXTERNAL**

WASH WITH WATER

**INTERNAL**

DRINK MUCILAGE, RAW EGG  
WHITE, MILK OR RICE GRUEL.  
FOLLOW WITH EMETIC. (TA-  
BLESPOONFUL OF MUS-  
TARD IN GLASS OF WATER).

**CALL A PHYSICIAN**

Do not mix Agway Sanitizer and  
Teat Dip with acid detergents.  
Use dilute solutions only as di-  
rected on label.

### STORAGE

Degrades with age. Use a test kit  
and increase dosage as necessary  
to obtain required level of avail-  
able chlorine.

Rinse empty container thoroughly  
with water and discard it.

## TABLE OF PROPORTIONS

Available Chlorine: 200 ppm— Use 1 oz. to 3 gallons of water.

## DIRECTIONS FOR SANITIZING PIPELINES, BULK TANKS, MILKING MACHINES, AND UTENSILS

After cleaning and just prior to use, sanitize all equipment with a solution of 200 ppm available chlorine. All surfaces should be exposed to sanitizer for a period of not less than two minutes. Do not rinse equipment with water after sanitizing. Do not soak overnight.

Keep in cool, dark place as light and heat will reduce the strength.

## DIRECTIONS FOR TEAT DIPPING

Use as a post-milking teat dip as an aid in reducing the spread of organisms which may cause mastitis.

Make fresh solution at each milking at a ratio of one (1) part Agway Sanitizer and Teat Dip to one (1) part of water, in a plastic pail or jug. Fill teat dip cup with solution and immediately after milking dip each teat up to base of the udder. Use a fresh cup of solution for each cow. Allow to air dry. Do not wipe.

