

TOXAPHENE 6E

NET CONTENTS
5 GALLONS

(Contains 6 pounds Toxaphene per gallon)

ACTIVE INGREDIENTS:	% by weight
*Toxaphene	60.0
Xylene	35.0
INERT INGREDIENTS:	5.0
	100.0

*Technical chlorinated camphene (67-69% chlorine content)

WARNING:

**KEEP OUT OF REACH OF CHILDREN
READ ENTIRE LABEL FOR ADDITIONAL HANDLING
PRECAUTIONS.**

May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe spray mist.

Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Do not use on cucurbits. This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Destroy all containers when empty by punching holes in them and burying with wastes. Do not burn. Do not reuse containers for other purposes. Rinse equipment and containers and dispose of wastes by burying in noncrop lands away from water supplies.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

ANTIDOTE: Internal: Give sodium chloride 15 g. (one table-spoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately.

External: Wash with soap and water.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

GRAIN CROPS:

CORN: (Preemergence or preplant broadcast only.) For control of armyworms and cutworms use 1½ to 2 quarts of this product suitably diluted with water per acre.

WHEAT, BARLEY, OATS, RYE, RICE, GRAIN SORGHUM: For control of armyworms and cutworms use 2⅔ pints of this product suitably diluted with water per acre.

Do not ensile treated sorghum forage.

Do not apply to sorghum within 40 days of harvest. Do not apply more than one treatment after heads start to form.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

CABBAGE: For control of cutworms (soil): Apply 2 to 3 pounds of active ingredient per acre, or 1⅓ to 2 quarts of this formulation, suitably diluted broadcast on the soil surface.

For control of cabbage looper and imported cabbageworm, use 2 to 3 pounds active per acre or 1⅓ to 2 quarts of this formulation.

DO NOT apply after heads start to form.

Plants treated with toxaphene must not be fed to dairy animals or animals being finished for slaughter.

COLLARDS AND KALE: For control of climbing and surface-feeding cutworms, thrips, serpentine leaf miner, armyworm, cabbageworm and green peach aphids, apply at the rate of 2⅔ pints per acre to give good distribution.

For control of cabbage looper, apply at the rate of 2⅔ to 5⅓ pints per acre in sufficient water to give good distribution. Do not make more than a total of 4 applications nor more than 3 applications after thinning. Do not apply the high rate within 35 days of harvest. Do not apply the low rate within 28 days of harvest.

SOYBEANS: For control of velvetbean caterpillar, bollworm, corn earworm, bean leaf beetles, armyworms, grasshoppers, blister beetle, green cloverworm, use 1½ to 2⅔ quarts of this material suitably diluted with water per acre. This product may be applied with either aerial or ground sprayers. With aerial application, the desired

volume of insecticide concentrate should be diluted with water and applied at 3 to 4 gallons per acre.

Avoid spray drift. Do not apply less than 3 weeks before harvest. Do not feed plants treated with toxaphene, or ensilage made from treated plants, to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

TOMATOES: For control of tomato hornworm, tomato fruitworm, Colorado potato beetle, pinworm, armyworm, cutworm, serpentine leaf miner, flea beetle and russet mite, apply the rate of from 2⅔ pints to 6⅔ pints (2 to 5 pounds of actual toxaphene) per acre, depending on the size of the plants. Do not apply within 3 days of harvest.

PEARS: For control of pear psylla, mites and red-banded leaf roller, apply from 1½ to 1⅔ pints of toxaphene emulsifiable concentrate per 100 gallons of spray. Apply as full coverage spray. Do not apply within 40 days of harvest or after second cover spray.

COTTON: For control of boll weevil, bollworm, cutworms, fall armyworm, lygus, leafworm, grasshoppers, use 2⅔ to 5⅓ pints of this material suitably diluted with water per acre. For control of thrips and fleahoppers use 1 pint suitably diluted in water per acre.

Do not graze nor feed treated cotton trash to dairy or animals being finished for slaughter.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

**Agway Inc., Fertilizer-Chemical Division,
Box 1333, Syracuse, New York 13201**