


4 lbs. NET WEIGHT



poultry shield

Toxi-FilmTM
Sevin* 50W

ACTIVE INGREDIENT:		
Carbaryl (1-naphthyl N-methylcarbamate)		50.00%
INERT INGREDIENTS:		50.00%
	Total	100.00%

For control of Poultry Pests on Chickens, Turkeys, Ducks, Geese, Game Birds and Pigeons.
Read Entire Label. Use Strictly in Accordance with Label Cautions, Warnings, and Directions.


FC21 D **CAUTION:** USDA Reg. No. 8590-21

**KEEP OUT OF REACH OF CHILDREN AND ANIMALS
HARMFUL IF INHALED OR SWALLOWED.**

(SEE ADDITIONAL CAUTIONS ON BACK PANEL)

Do not breathe the dust or spray mist. Do not take internally. SKIN CONTACT MAY BE IRRITATING. Avoid contact. Wash hands before eating. Take shower or bath after use. WEAR REGULAR LONG SLEEVED WORK CLOTHING. Change to clean clothing daily. Avoid contamination of feed and foodstuff. Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

*Trademark of Union Carbide Corporation for the active ingredient Carbaryl (1-naphthyl N-methylcarbamate)



WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Caution: Do not apply this material within 7 days of slaughter either to the birds or to quarters. Avoid contamination of nests, eggs, feeding and watering troughs.

Direct Mist Spray on Birds: Control northern fowl mite, chicken mite, lice and fleas by

- (1) Misting with clip-on fogger: Mix 10 ounces of Sevin 50W in 1 gallon of spray. Use 1 1/2 gallons per 1000 hens in cages, on litter or on slatted floor. Repeat in 4 weeks if necessary, or
- (2) Spraying with knapsack or cylinder type compressed air sprayers: Mix 6 ounces of Sevin 50W in 5 gallons of spray. Use 1 gallon per 100 hens in cages on litter or on slatted floor. Repeat in 4 weeks if necessary.

Direct mist spraying for chicken mites and fleas is a supplement to spraying roosts and buildings for control of these pests.

Roosts and Buildings: Spray with conventional power spray or knapsack equipment. For chicken mites, fleas and bedbugs, use 2 pounds per 25 gallons of water. For fowl ticks, use 8 pounds per 25 gallons of water. Spray 1 to 2 gallons per 1000 square feet of wall, bedding, litter or roost surfaces. Force spray into cracks and crevices. Repeat as needed. Ventilate while spraying.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

MANUFACTURED BY AGWAY INC. - Chemical Division, Box 1333, Syracuse, N.Y. 13201

