

# FERBAM 7.6 D

NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENT:

Ferbam (ferric dimethyl dithiocarbamate) ..... 7.6%

INERT INGREDIENTS: ..... 92.4%

100.0%

## CAUTION:


KEEP OUT OF REACH OF CHILDREN.

May cause irritation of nose, throat, and skin. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. This product is toxic to fish and wildlife. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

## DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.



Consult state agricultural extension service for information on exact timing of applications on the crops mentioned.

**APPLES:** Scab, Cedar-Apple Rust, Black Rot, Blotch, Bitter Rot, Brook's Spot: Use 1 to 2 lbs. per average size tree. Apply in pre-bloom, calyx, and cover periods. Do not apply within 7 days of harvest.

**PEARS:** Scab, Sooty Blotch, Fabraea Leaf Blight, Fruit Spot: Use 1 lb. per average size tree. Apply in pink, calyx, and cover periods. Do not apply within 7 days of harvest.

**CHERRIES:** Brown Rot: Use 1 lb. per average size tree. Apply in popcorn, full-bloom, petal-fall, and cover periods.

**PEACHES:** Brown Rot Blossom Blight: Use 1 lb. per tree. Apply in pink bud and 25% to 75% bloom periods. Do not apply within 21 days of harvest.

**BLUEBERRIES:** For control of red leaf disease and blossom blight apply 15 to 20 lbs. per acre. On new burn, make application in late June or early July when fungus appears on under-side of leaves on bearing bushes. On bearing bushes, make first application when buds swell, repeat when blossoms first open and again during extended wet periods. Apply also in late June or early July for red leaf disease. Do not use within 40 days of harvest.

**TOBACCO:** Blue Mold in plant beds: Make first application when leaves are size of a dime; use 1 to 1½ lbs. per 100 sq. yds. Continue applications twice weekly, increase dosage gradually up to 3½ lbs. per 100 sq. yds. when plants are ready for field-setting.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Code No. AC32E

USDA Reg. No. 8590-32

Manufactured by Agway Inc., Syracuse, N. Y.