

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 4590-2



# CAPTAN-PARATHION 10-2 D

ACTIVE INGREDIENTS:

Captan (N-trichloro-methylmercapto-4-cyclohexene-1, 2-dicarboximide) .....	10.0%
Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate) .....	2.0%
INERT INGREDIENTS: .....	88.0%
	<hr/>
	100.0%



**DANGER:**  
**KEEP OUT OF REACH OF CHILDREN.**

Poisonous if swallowed, inhaled, or absorbed through skin. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for parathion protection. Rapidly absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing, and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep out of reach of children. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. During application avoid drift to adjacent food and forage plantings which might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse this empty container. Dispose of by burying with wastes or burning. Stay away from smoke and fumes.

ANTIDOTES: If swallowed: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down

and keep quiet. CALL A PHYSICIAN IMMEDIATELY! If on skin: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

## DIRECTIONS

**READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.**

Apples: To aid in control of scab, Brook's fruit spot, sooty blotch, fly speck, codling moth, plum curculio, leaf rollers, bud moth, redbug, aphids and mites, apply at the rate of 30-50 pounds per acre using the higher rate to control bud moth and redbug. Full-coverage applications by airplane or ground duster is necessary for satisfactory control.

Repeat application at 4 to 7 day intervals as needed to maintain cover when apple scab is prevalent in the pre-bloom, bloom, petal-fall, first and second cover dust application. Continue cover sprays at 7 to 14 day intervals during growing season as necessary for disease and insect control. Do not apply during blossom period.

**CAUTION:** Do not apply within 14 days of harvest. Remove residues remaining on fruit at harvest by washing, wiping, or other effective means. Do not apply within 10 days of an oil spray. Use cautiously on Red Delicious and Baldwin varieties.

**NOTE:** This pesticide is to be sold ONLY in this original unbroken package.

Agway Inc., Chemical Division  
Box 1333, Syracuse, N. Y. 13201

EPA Reg. No. 8590-2



# MANEB-SEVIN 4.5-5 D

TO CONTROL CERTAIN INSECTS AND DISEASES OF VEGETABLES LISTED ON THIS LABEL

## ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate).....	5.0%
Maneb (manganese ethylene bisdithiocarbamate).....	4.5%
(Metallic manganese content 0.98%)	

## INERT INGREDIENTS

(not more than).....	90.5%
	<hr/> 100.0%

## DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Maneb-Sevin Dust has been formulated for control of both insects and diseases. Best results are obtained when the dust is applied before serious build-up of insects or disease occurs. Repeat application at 7 to 10 day intervals, making thorough coverage of top and underside of foliage.

**BEANS—LIMA, SNAP, DRY:** For control of Downy mildew, Mexican bean beetle, Lygus and Stink bugs, apply 20 lbs. dust per acre. Apply at 7 day intervals. Do not apply within 4 days of harvest on fresh snap beans.

**CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, KOHLRABI:** To control Downy mildew, Alternaria, Flea beetle and Harlequin bug, apply 20-30 lbs. dust per acre. Do not apply within 7 days of harvest on cabbage. Do not apply to cauliflower, broccoli, brussels sprouts, and kohlrabi within 3 days of harvest.

**CARROTS:** For control of Alternaria leaf blight, Cercospora leaf spot, Flea beetles and Leafhoppers, apply 20-30 lbs. dust per acre. To control six-spotted leafhoppers (Aster yellow vector), apply 30 lbs. dust per acre. Apply at 5 to 7 day intervals. Do not apply within 7 days of harvest. Do not use tops for food or feed.

**CUCUMBERS, CANTALOUPE, MELONS, SQUASH, WATERMELONS:** For the control of Anthracnose, Downy mildew, Cucumber beetle, Melonworm, Pickleworm, Squash Bugs and Flea beetles, apply 26 lbs. dust per acre. Avoid excessive applications. Some leaf injury

SP 774

may occur if treatments are made when tender foliage is wet or in presence of very high humidity. Do not apply within 5 days of harvest.

**LETTUCE, ENDIVE:** For control of Downy mildew, Flea beetle, Harlequin bug, Leafhoppers, apply 30 lbs. dust per acre. Apply at 7 day intervals. Do not apply within 10 days of harvest for head lettuce or 14 days of harvest for leaf lettuce and endive. Remove residue from head lettuce by stripping and trimming and from leaf lettuce and endive by washing or other effective means.

**POTATOES:** For control of early and late Blight, Colorado potato beetle, Leafhoppers, Flea Beetle, apply 30 lbs. dust per acre. As plants become larger and late Blight threatens, apply 40 lbs. dust per acre. Apply at 7 day intervals. If late Blight is present, shorten interval to 3 to 5 days. Foliage application only.

**SPINACH:** For control of Downy mildew, Flea beetle, Leafhopper, apply 30 lbs. dust per acre. Repeat at 7 to 10 day intervals. Do not dust within 14 days of harvest, and wash thoroughly before use.

**SWEET CORN:** For control of Helminthosporium leaf blight, Flea Beetle, Corn earworm, European corn borer, Fall armyworm, Sap beetles, Japanese beetle, apply 20 to 40 lbs. dust per acre. Apply at 3 day intervals. For larvae in the whorl and foliage feeders, apply to entire plant, repeat as necessary. For insect attacking silks and ears, apply at 2-3 day intervals starting when silks first appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestations. Timing and good coverage is absolutely essential for effective control. Do not feed treated forage to livestock. Application of the product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

**TOMATOES, EGGPLANT, PEPPERS:** For control of early and late Blight, Anthracnose, Stemphylium (Gray leaf spot), Septoria leaf spots, Tomato fruitworm, Fall armyworm, Flea beetles, Colorado potato beetle, European corn borer and Stink bug, apply 30 lbs. dust per acre. As plants become larger or late Blight threatens, apply 40 lbs. of dust per acre. Apply at 7 day intervals. If late Blight is present, shorten interval to 3 to 5 days. Do not apply within 5 days of harvest on tomatoes.

NET WEIGHT FIFTY POUNDS

To avoid damage do not use on young tender plants in the greenhouse.

**CAUTION:** KEEP OUT OF REACH OF CHILDREN

ACCEPTED  
MAR 27 1975  
EPA Reg. No. 8590-33

Do Not Take Internally. Avoid Breathing of Dust. Harmful if inhaled or swallowed. Skin Contact May Be Harmful. Avoid Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Do Not Contaminate or Store Near Feed or Food-stuffs.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

This product is toxic to fish. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

**NOTE:** Sevin contained in the dust is toxic to Honeybees. Avoid using this dust during flowering and pollination period.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

**NOTE:** This pesticide is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590-33 EPA Est. No. 8590-PA-1

**AGWAY INC.**  
**FERTILIZER — CHEMICAL DIVISION**  
**BOX 1333, SYRACUSE, N. Y. 13201**