

MANEB 4.5 D

FOR USE IN CONTROLLING CERTAIN DISEASES OF CROPS LISTED ON THIS LABEL

ACTIVE INGREDIENTS:

Maneb (manganese ethylene
bisdithiocarbamate) 4.5%

~~Diatomic manganese~~
~~Inert ingredients~~
Inert ingredients
(not more than) 95.5%

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS

Maneb 4.5 D can be used for the control of many common diseases of vegetables and field crops. Maneb may be applied to either moist or dry foliage any time dusting conditions are favorable. Applications should begin when plants are small. Use 25 to 50 pounds per acre depending on the size of plants. Repeat at 5 to 10 day intervals depending on the severity of disease conditions. Dusters should be adjusted to give uniform and thorough coverage.

TOMATOES: ~~Early and late blight~~, Anthracnose, Stemphylium (gray leaf spot), Septoria leaf spots — begin when disease is first reported in area and repeat applications at 7 to 10 day intervals or oftener under severe late blight conditions. Do not use on tender young plants in the greenhouse. Do not apply within 5 days of harvest.

POTATOES: ~~Early and late blight~~—apply every 5 to 10 days throughout the season. More often under severe late blight conditions.

Begin applications at 15 to 20 lbs. per acre and increase poundage as necessary to provide adequate coverage as vines increase in size.

CELERY: ~~Early and late blight~~—begin dusting in the plant bed when plants first come through the ground. Make further applications at 3 to 5 day intervals until plants are set in the field. Continue field applications on regular weekly schedule. Remove residue by stripping, trimming and washing. Do not apply within 14 days of harvest.

CUCUMBERS, CANTALOUPE, MELONS, SQUASH, WATERMELONS: ~~Anthracnose and Downy mildew~~—start applications when runners begin to form or at the first sign of disease. Repeat at weekly intervals. Direct dust to underside of leaves. Do not apply within 5 days of harvest.

LIMA BEANS, DRY BEANS, SNAP BEANS: ~~Downy mildew~~—apply when mildew first appears and repeat at weekly intervals. Do not use within 4 days of harvest on fresh snap beans.

ONIONS: ~~Downy mildew, Blast, Botrytis leaf blight~~ Purple blotch—begin applications when plants reach 4 leaf stage or when diseases are reported in the area and continue at 7 to 10 day intervals during the season. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.

SPINACH: ~~Downy mildew (bacterial)~~—apply at first sign of mildew and repeat at 7 to 10 day intervals or as required for adequate control. Do not dust within 10 days of harvest, and wash thoroughly before use.

SWEET CORN: ~~Helminthosporium leaf blight~~—repeat applications at 3 to 7 day intervals as long as disease threatens. Do not feed treated forage to livestock.

CAUTION: KEEP OUT OF REACH OF CHILDREN:

May cause irritation of nose, throat, eyes and skin. Avoid inhalation of dust. In case of skin contact wash thoroughly with water. Store in cool dry place. Keep container closed when not in use.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. Do not reuse empty container. Destroy it by burying wastes or by burning. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590-42 EPA Est. No. 8590-PA-1
AGWAY INC.

**FERTILIZER — CHEMICAL DIVISION
BOX 1333, SYRACUSE, N. Y. 13201**