

ADVERTISING AND SALES PROMOTION SERVICES

303W

FOR: MANEB 4.5D

DATE: 1/5/83

((FRONT PANEL))

NET WEIGHT

50#

AGWAY

MANEB 4.5D

ACTIVE INGREDIENT:

Maneb (Manganese ethylene bisdithio-
carbamate)

4.5%

(Metallic manganese
content 0.98%)

INERT INGREDIENT

(not more than)


95.5%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)

See additional precautionary statements on side
panel.



AGWAY INC.

CHEMICAL DIVISION

BOX 4741

SYRACUSE, NY 13221

OK BY

EPA Reg. No. 8590-42

EPA Est. No. 8590-PA-1

SHEET NO 1 - 7

ADVERTISING AND SALES PROMOTION SERVICES

303W

FOR:

MANEB 4.5D

DATE: 1/5/83

((REAR PANEL))

DIRECTIONS FOR USE

It is a violation to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL (12 pt.)

Do not contaminate water, food or feed by storage or disposal.

Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

Completely empty bag by shaking & tapping sides & bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State & local authorities.

READ THESE ENTIRE DIRECTIONS FOR USE AND USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS & DIRECTIONS FOR USE.

O'K BY

SHEET NO 2 - 7

ADVERTISING AND SALES PROMOTION SERVICES

303Y

FOR:

MANEB 4.5D

DATE:1/5/83

Maneb 4.5D can be used for the control of many common diseases of vegetables and field crops. Maneb may be applied to either moist or dry foliage any time dusting conditions are favorable. Applications should begin when plants are small. Use 25 to 50 pounds per acre depending on the size of plants. Repeat at 5 to 10 day intervals depending on the severity of disease conditions. Dusters should be adjusted to give uniform and thorough coverage.

TOMATOES: Early and late blight, Anthracnose, Stemphylium (gray leaf spot), Septoria leaf spots - begin when disease is first reported in area and repeat applications at 7 to 10 day intervals or oftener under severe late blight conditions. Do not use on tender young plants in the greenhouse. Do not apply within 5 days of harvest.

POTATOES: Early and late blight - apply every 5 to 10 days throughout the season. More often under severe late blight conditions. Begin applications at 15 to 20 lbs. per acre and increase poundage as necessary to provide adequate coverage as vines increase in size.

OKBY

SHEET NO 3 - 7

ADVERTISING AND SALES PROMOTION SERVICES

3030

FOR:

MANEB 4.5D

DATE: 1/5/83

CELERY: Early and late blight - begin dusting in the plant bed when plants first come through the ground. Make further applications at 3 to 5 day intervals until plants are set in the field. Continue field applications on regular weekly schedule. Remove residue by stripping, trimming and washing. Do not apply within 14 days of harvest.

CARROTS: Alternaria leaf blight, Cercospora leaf spot - begin applications when diseases are reported in the area and continue at 7 to 10 day intervals during the growing season, or as required for adequate control.

CUCUMBERS, CANTALOUPE, MELONS, SQUASH, WATER-MELONS: Anthracnose and Downy mildew - start applications when runners begin to form or at the first sign of disease. Repeat at weekly intervals. Direct dust to underside of leaves. Do not apply within 5 days of harvest.

LIMA BEANS, DRY BEANS, SNAP BEANS: Downy mildew - apply when mildew first appears and repeat at weekly intervals. Do not use within 4 days of harvest on fresh snap beans.

O.K BY

SHEET NO 4 - 7

ADVERTISING AND SALES PROMOTION SERVICES

3030

FOR:

MANEB 4.5D

DATE: 1/5/83

ONIONS: Downy mildew, Blast, Botrytis leaf blight Purple blotch - begin applications when plants reach 4 leaf stage or when diseases are reported in the area and continue at 7 to 10 day intervals during the season. Do not apply within 7 days of harvest. Do not apply to exposed bulbs.

SPINACH: Downy mildew (bluemold) - apply at first sign of mildew and repeat at 7 to 10 day intervals or as required for adequate control. Do not dust within 10 days of harvest, and wash thoroughly before use.

SWEET CORN: Helminthosporium leaf blights - repeat applications at 3 to 7 day intervals as long as disease threatens. Do not feed treated forage to livestock.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention.

OK BY

SHEET NO 5 - 7

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

MANEB 4.5D

DATE: 1/5/83

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

If on skin, remove by washing.

If in eyes, get medical attention.

Wear protective clothing (long pants, long sleeve shirt, gloves, hat & boots) during mixing and loading.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from fire and sparks. Store in a cool, dry place.

OK BY

SHEET NO 6 - 7

ADVERTISING AND SALES PROMOTION SERVICES

309V

FOR:

MANEB 4.5D

DATE:1/5/83

AGWAY DISCLAIMS ALL IMPLIED WARRANTIES, INCLUDING IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL INJURY, LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR DAMAGES, NOT TO EXCEED THE PURCHASE PRICE OF THE PRODUCT AND SHALL NOT INCLUDE ANY INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

O'K BY

SHEET NO 7 - 7