



SEVIN 3 D

NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENTS:

*Carbaryl (1-Naphthyl N-methylcarbamate) 3.0%

INERT INGREDIENTS,
not more than 97.0%
100.0%

*Trademark of Union Carbide Corp. for the active ingredient 1-Naphthyl N-methylcarbamate and is covered by U.S. Patent No. 2,903,478.

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Harmful if Swallowed. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. Skin contact with dry insecticide may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is toxic to fish and wildlife. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty

of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Thorough coverage of entire plant, including lower leaf surface, is essential for best insect control. Do not use with lime or other highly alkaline materials. Suitable for aircraft application.

BEANS: To control Mexican bean beetle, apply 20 lbs. per acre. For leafhoppers, cucumber beetles, flea beetles and Japanese beetles, apply 35 lbs. per acre. For armyworms, corn earworm and velvet bean caterpillars, apply 50 lbs. per acre. For lima bean pod borer, stink bugs and tarnished plant bugs, apply 65 lbs. per acre. Apply when insects appear. Repeat as needed. May be applied up to harvest.

CORN: To control corn earworm, begin application depending on moth flight and egg laying. Repeat every 2-3 days. Apply as needed to control fall armyworm and Japanese beetle. Use 50 lbs. per acre. May be applied up to harvest.

CUCUMBER & SUMMER SQUASH: To control flea beetles, melonworms, pickleworms, squash bugs and striped and spotted cucumber beetles, apply 30 lbs. per acre when insects first appear. Repeat at 7 to 10 day intervals as needed. May be applied up to harvest.

EGGPLANT, PEPPERS & TOMATOES: To control fall armyworm, flea beetles, leafhoppers, and tomato fruitworm, apply 35 lbs. per acre. For European corn borer, stink bugs and tomato hornworm, apply 50 lbs. per acre. Apply when insects first appear. Repeat at 7-10 day intervals as needed. May be applied up to harvest.

POTATOES: To control Colorado potato beetle apply 25 lbs. per acre. For leafhoppers and flea beetles, apply 35 lbs. per acre. For tarnished plant bugs, apply 60 lbs. per acre. May be applied up to harvest. Foliage application only.

TIMING OF APPLICATION: Timing and rate of application vary as a result of local conditions. Since these conditions are unknown to the manufacturer, you must satisfy yourself as to local requirements. Do not exceed maximum rates or use closer to harvest than specified above. Avoid contamination of feed and foodstuffs.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Code No. AC43E
USDA Reg. No. 8590-43

Manufactured by Agway Inc., Syracuse, N. Y.

