

DIRECTIONS FOR USE (Continued)

PUMPKINS AND SQUASH: For control of Cucumber Beetle, Squash Bug and Squash Vine Borer. Apply 1 to 1½ pints. Do not apply after edible portions begin to form. There may be occasional plant injury.

SWEET POTATO: For control of Flea Beetle Larvae, White Grubs, Wireworms. Apply 2½ to 5 pints, before planting and work into the top few inches of soil.

TOMATO: For control of Flea Beetles, Serpentine Leaf-miner and Tomato Hornworm. Apply 1 to 2 pints. Several repeat applications may be necessary. Do not apply after the fruit begins to form.

WATERMELON: For control of Cucumber Beetles, Squash Bug, and Squash Vine Borer. Apply 1 to 1½ pints. Do not apply after edible portions begin to form.

FRUITS

GENERAL: Apply in 100 gallons of water per acre unless otherwise directed.

BLACKBERRIES: For control of Strawberry Root Weevil (Adults). Apply 2 pints to the foliage. Apply before fruits appear in the Spring and after harvest. For control of White Grubs, Root Weevil Larvae. Apply 10 pints as a soil treatment. Work into the soil at the base of the plant when no fruit is present.

BLUEBERRIES: For control of Cranberry Rootworm and White Grubs. Apply 10 pints to the soil around the base of the plants when no fruit is present.

STRAWBERRIES: For control of Strawberry Root Weevil, Strawberry Rootworm and White Grubs. Apply 10 pints. Work into the soil prior to planting.

FIELD CROPS

GENERAL: Apply in 100 gallons of water per acre unless otherwise directed.

CORN (FIELD AND SWEET): For control of Asiatic Garden Beetles, Cutworms, Green June Bug Larvae, Japanese Beetles, Seed Corn Beetles, Seed Corn Maggots, Wireworms, White-Fringed Beetles and White Grubs. Apply 2½ to 4 pints as a broadcast treatment after field is plowed and prior to planting. Disc in promptly. For control of Corn Rootworms, Northern Corn Rootworms, Southern Corn Rootworms, Cornfield Ants, Corn Root Aphids. Apply 2 to 4 pints
SP868

broadcast or one pint in the row. As a broadcast treatment, apply after the field is plowed. Disc in promptly. Apply in the row at planting. For control of White Grubs and Wireworms. Apply 2 pints in the row. Apply at planting time.

OATS, RYE, BARLEY AND WHEAT: For control of Armyworms. Apply 1 to 2 pints when larvae are present. Do not apply after heads start to form. Do not graze treated foliage.

TOBACCO: For control of Wireworms. In transplanting water, add 2 oz. per 50 gallons of water for protecting the transplant seedlings. Use 250-300 gallons per acre. 4 ounces of Chlordane per 50 gallons of water is desired for heavy infestations.

SUBTERRANEAN TERMITE CONTROL: Use 1 quart of Chlordane per 50 gallons of water (4 teaspoonsful/gallon). Dig a 1 foot x 1 foot trench along exterior of foundation. Treat both trench and back fill. Apply ½ gallon of spray per linear foot of trench.

For deep foundations bore 1 inch holes 8 inches apart and 15 inches deep in the bottom of the trench. (Do not dig below the depth of footings.) Spray trench and back fill with 1 gallon of above spray per linear foot of trench.

Saturate unexposed soil areas under patios and walks adjacent to the foundation with the above spray. Pour in expansion cracks and in shallow trenches or holes bored 12 inches apart along walk edge or patio.

Treat exposed soil under porches with ½ gallon of same spray per square yard. In heavy clay soils increase the mixture to 2 quarts of Chlordane per 50 gallons of water (8 teaspoonsful/gallon). Spray thoroughly all areas where wood contacts soil as well as points of entry around basement windows, and pipes. Use 2 quarts per 50 gallons of spray (8 teaspoonsful/gallon).

Inspect annually to determine the need for reapplication.

HOUSEHOLD INSECTS: To control Ants, Box Elder Bugs, Carpet Beetles, Cockroaches, Crickets, Earwigs, Silverfish, Spiders and Wasps, use 5 table-spoonsful Chlordane per gallon of water. Apply as a coarse spray or paint infested areas paying special attention to baseboards, cracks, underneath rugs or other hiding places. Also treat doorways and other entry points. Repeat as needed. Use care in treating finished surfaces to avoid staining.



Agway

TM

NET CONTENTS ONE GALL

CHLORDANE 8

ACTIVE INGREDIENTS:	
Technical Chlordane*	72.0%
Petroleum Distillate	18.0%
INERT INGREDIENTS	
TOTAL	100.0%

*Equivalent to 43.2% Octachloro-4, 7-Methanotetra-dane and 28.8% related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Contact with skin can cause symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid ingestion of feed and foodstuffs.

This product is toxic to fish and wildlife. Keep out of streams and ponds. Birds feeding on treated areas killed. Apply this product only as specified on the label. Rinse equipment and containers and dispose of water by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in the bottom and burying with wastes, or by burning (keep out of smoke).

ALLON

8 E

72.0% by wt.
18.0% by wt.
10.0% by wt.
100.0% bywt.
notetrahydroin-

REN.

n cause toxic
case of con-
Avoid contam-

out of lakes,
areas may be
on this label.
of wastes by
supplies. Con-
; in them and
if smoke).

DIRECTIONS FOR USE

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

VEGETABLE CROPS

GENERAL: Apply in 100 gallons of water per acre unless otherwise directed.

ASPARAGUS: For control of Asparagus Beetle. Apply ½ pint. Make post cutting applications only.

BEANS AND BLACK-EYED PEAS: For control of Flea Beetles and Green Stink Bugs. Apply one to two pints. Do not make application to green snap beans after the pods begin to form.

CABBAGE AND OTHER COLE CROPS (BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KALE AND KOHLRABI): For control of Cabbage Maggot. Apply one pint. Newly set plants: Spray around base of plants as soon as set in the field. Stir or wash the treatment into the soil. One or two repeated applications are frequently necessary. For Transplanting Dip add two quarts per 100 gallons of water. Swish roots and crown through mixture just before setting. Protection is limited to newly set plants. Transplant drench: Add ¼ pint Chlordane per 50 gallons of water and apply around plants at planting time. Use about 400 gallons per acre. Protection is limited to newly set plants. For control of Cabbage Worms, Flea Beetles, Green Stink Bugs, and Harlequin Bugs. Apply one to two pints in 100 gallons. Several repeat applications may be necessary. No application after the heads or other edible portions begin to form.

CANTALOUPE, MELONS AND CUCUMBERS: For control of Cucumber Beetle, Squash Bugs, Squash Vine Borers. Apply one to one and one-half pints. There may be occasional slight plant injury. Do not apply after edible portions begin to form.

CARROTS: For control of Carrot Rust Fly. Apply one to two pints in 100 gallons. Several repeat applications would be advised. Treated tops should not be used as food.

EGGPLANTS: For control of Flea Beetle and Serpentine Leafminer. Apply one to two pints. Make repeated spray applications as necessary. There should be no application after the fruit begins forming.

LETTUCE: For control of Armyworm, Lygus Bug, Serpentine Leafminer, Spotted Cucumber Beetle. Apply one and one-half to two pints. Do not make application after heads begin to form on head lettuce or after seedling stage on leaf lettuce.

ONIONS: For control of Onion Maggot. Apply one pint. Spray around the base of plants at transplanting time only. Do not use on green or Spring onions. For control of thrips. Apply 1½ pints. Do not make applications to green or Spring onions.

PEAS: For control of Pea Leafminer. Apply 1 to 1½ pints. Repeat applications as necessary. Do not feed treated pea vine to dairy animals or animals being fattened for slaughter. For blackeyed peas do not apply after the pods begin to form.

PEPPERS: For control of Cutworms. Apply 1½ to 2 pints. Work into the soil around the plant at transplanting time only. Do not make application after the peppers begin to form. For control of Mole Crickets. Apply 1 pint at transplanting time only. Work into the soil around the base of the plant. Do not make application after peppers begin to form. For control of Wireworm. Apply 2½ to 5 pints. Work into soil prior to or at planting in the field.

POTATOES: For control of Wireworms. Apply 2½ to 6 pints. Work into the soil prior to or at planting time. Applications of 5 to 10 pints will give three years control. If application is made to the row use ¼ to ½ of the dosage for broadcast application.