

# COPPER 7 D

## NET WEIGHT FIFTY POUNDS

### ACTIVE INGREDIENT:

Copper (in basic copper sulfate) expressed as  
metallic copper ..... 7.0%

### INERT INGREDIENTS,

not more than ..... 93.0%  
100.0%


## CAUTION:

**KEEP OUT OF REACH OF CHILDREN. HARMFUL IF  
TAKEN INTERNALLY.**

This product is toxic to fish. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

SP 168



## DIRECTIONS

**READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.**

For best disease control, dust all plants thoroughly, particularly the undersides of the leaves. If insect control on these crops is needed, use other Dusts containing Metallic Copper plus the proper insecticides.

**CUCUMBERS, MELONS AND RELATED CROPS:** To prevent downy mildew and angular leaf spot, apply 45 to 55 lbs. per acre at 5 to 7 day intervals whenever weather conditions are conducive to the development of the fungus. Do not apply to young melons before runners are formed.

**POTATOES AND CELERY:** To prevent <sup>blight</sup> early and late blights, begin applications when plants emerge. Apply at weekly intervals throughout season at 30 to 50 lbs. per acre, depending on size of plants.

**TOMATOES:** To control bacterial spot, use 30 to 40 lbs. per acre at weekly intervals in early flowering stage. When late blight threatens, apply at rate of 50 lbs. per acre.

**TIMING OF APPLICATION:** Timing and rate of application vary as a result of local conditions. Since these conditions are unknown to the manufacturer, you must satisfy yourself as to local requirements. Your state agricultural extension service may be able to give you information.

**NOTE:** This pesticide is to be sold ONLY in this original unbroken package.

Code No. AC51E

USDA Reg. No. 8590-51

Manufactured by Agway Inc., Syracuse, N. Y.