



NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENTS:

Captan (Methylcaptafol
1,2-dichlorophenyl
ethyl dimethylcarbamate)

7.5%

INERT INGREDIENTS:

92.5%

100.0%

STORE IN COOL, DRY PLACE PROTECT FROM EXCESSIVE HEAT

CAUTION:

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Avoid inhalation of dust, avoid prolonged or repeated contact with skin. Avoid contamination of feed and food-stuffs.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

SP-547

DIRECTIONS

USE ONLY AS DIRECTED AND STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS. KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.

Captan Dust is a fungicidal dust formulation suitable for use by ground duster or airplane.

APPLES: Apple Scab, Brooks Fruit Spot, Fire Spotting, Bitter Rot, Black Rot, Botrytis, 30 to 35 lbs. per acre. Make first application as soon as spore spores are ready to discharge and an infection period threatens. Repeat application for each subsequent infection period.

CELERY: (Field and Seed Bed Only): Pink Rot, Damp-off Disease. Use 25-30 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

CANTALOUPES, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS: Downy Mildew, Angular Leaf Spot, Anthracnose. Apply at 25 to 30 lbs. per acre on young plants, increasing to 35 lbs. per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 10 day intervals until danger of infection is past.

ORNAMENTALS: ROSES, Black Spot; CARNATIONS, Rust and Leaf Spot; CHRYSANTHEMUMS, Flower Blight. Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

PEACH, NECTARINES: Brown Rot and Peach Scab. 80 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal fall and 3 to 4 weeks as necessary. Make pre-harvest application where necessary. Do not apply within 1 day of harvest.

POTATOES (Seed Pieces): Seed Rot and Late Blight. Use at 1 lb. to 1 1/2 lbs. per seed piece. Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging.

SOIL AND GREENHOUSE BENCH TREATMENT: Preventing for Damp-off and Root Rot Diseases on Seeding or Transplants of Roots (and other shrubs and trees). Many flowers and vegetables (such as Beans, Peas, Cucurbits) and lawn seedbeds—1 1/2 to 2 1/4 lbs. per 1000 sq. ft. (50 to 60 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

TIMING OF APPLICATION: Timing and rate of application vary as a result of local conditions. Since these conditions are unknown to the manufacturer, you must satisfy yourself as to local requirements. Do not exceed maximum rates or use closer to harvest than specified above.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Code No. AC52E

USDA Reg. No. 8590-52

Manufactured by Agway Inc., Syracuse, N. Y.

