AGWAY CAPTAN 7.5D

FOR USE IN CONTROLLING FRUIT, VEGETABLE AND ORNAMENTAL DISEASES.

ACTIVE INGREDIENTS:

Captan (N-trichloromethyl-thio-4cyclohexene-1, 2-dicarboximide 7.10%
Related Compounds 0.40%
INERT INGREDIENT: 92.50%

ACCEPTION 4 1981

TOTAL
KEEP OUT OF REA TH OF CHILDREN.

CAU ION

See Additional Cautions Below.

STORE IN COOL DRY PLACE, PROTECT FROM EXCESSIVE HEAT.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Causes skin and eye irritation. May be harmful if swallowed or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Use with adequate ventilation. Wash thoroughly after handling.

FIRST AID: In case of eye contact, immediately flush eyes with plenty of water for at least 15 minutes. Call a physician. Wash skin with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Captan 7.5D is a fungicidal dust formulation suitable for use by ground duster or airplane.

APPLES: Apple Scab, Brooks Fruit Spot, Fly Speckling, Bitter Rot, Black Rot, Botryosphaeria. 30 to 35 lbs. per acre. Make first application as soon as scab spores are ready to discharge and an infection period threatens. Repeat application for each subsequent infection period.

CELERY: (Field and Seed Bed Only): Pink Rot, Damp-off Disease. Use 25-30 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

CANTALOUPES, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS: Downy Mildew, Angular Leaf Spot, Anthracnose. Apply at 25 to 30 lbs. per acre on young plants, increasing to 35 lbs per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 10 day intervals until danger of infection is past.

ORNAMENTALS: ROSES, Black Spct; CARNATIONS, Rust and Leaf Spots; CHRYSANTHEMUMS, Flower Blight. Apply at first sign of disease to give evan coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

PEACH, NECTARINES: Brown Rot and Peach Scab. 50 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary. Do not apply within one day of harvest.

100.00%

POTATOES (Seed Piaces): Seed Rot and Damp-off. Dust at 1 lb. to 120 lbs. cut seed pieces. Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within six hours of cutting. If planting is to be delayed more than one or two days, the treated pieces should be stored for two days in open crates before bagging.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for Damp-off and Root Rot Diseases on Seedling or Transplanting of Roses (and other shrubs and trees), many flowers and vegetables (such as Beans, Peas, Cucurbits) and lawn seedbeds — 1½ to 2½ lbs. per 1000sq. ft. (50 to 60 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

TIMING OF APPLICATION: Timing and rate of application vary as a result of local conditions. Since these conditions are unknown to the manufacturer, you must satisfy yourself as to local requirements. Do not exceed maximum rates or use closer to harvest than specified above.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not is strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Pederal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bag according to approved Federal, State or local procedures under the Resource Conservation and Recovery. Act

EPA Reg Na 8590-52

Era Est No 8590-PA-1

AGWAY INC., CHEMICAL DIVISION BOX 4933, SYRACUSE, NY 13221

Net Contents 50 lbs.