

NET CONTENTS

AGWAY

FIFTY POUNDS

METHOXYCHLOR 5 D

ACTIVE INGREDIENTS:

* Methoxychlor, technical.....	5.0%
INERT INGREDIENTS, not more than.....	95.0%
	100.0%

*Equivalent to 4.4% 2, 2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 0.6% other isomers and related compounds.

CAUTION KEEP OUT OF REACH OF CHILDREN.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of dust from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning (keep out of smoke). This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Thorough application to all plants is essential for adequate insect control.

BEANS: To control both Mexican bean beetle and leafhoppers, use 25 to 35 lbs. per acre, depending on size of plants. Apply when insects first appear and repeat at 5 to 7 day intervals as necessary. Do not apply within 3 days of harvest. Do not feed treated vines to livestock.

CUCUMBERS, MELONS AND SUMMER AND WINTER SQUASH: To control aphids, cucumber beetles, flea beetles, leafhoppers, squash bugs and squash vine borers, use 35 lbs. per acre. Apply when insects first appear and repeat as necessary. Do not apply within 1 day of harvest.

ALFALFA AND CLOVER: To aid in the control of alfalfa weevil, meadow spittlebug, and potato leafhoppers, use 30 lbs. per acre. Apply when insects first appear. Do not apply within 7 days of harvest. Do not apply more than one application. If possible, cut 7 to 10 days following treatment.

GRAPES: To control grape berry moth, use 60 lbs. per acre. Make first application just after bloom, second application 10 days later, and third application when brood berry moth appears. Do not apply within 14 days of harvest. Use this material in combination with other Special Grape Dusts containing copper, ferbam, and an insecticide with copper or ferbam for a complete grape dust program.

BLUEBERRIES: For control of Japanese beetle, Blueberry maggot, leafhoppers, and spittlebug. Apply 25 to 35 lbs. per acre starting when pests first appear and continuing at 7-10 day intervals or as needed. Do not apply within 14 days of harvest.

TIMING OF APPLICATION: Timing and rate of application vary as a result of local conditions. Since these conditions are unknown to the manufacturer, consult your State Agricultural Extension Service for local requirements. Do not exceed maximum rates or use closer to harvest than specified above.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Manufactured by Agway Inc., Chemical Division, Box 1333, Syracuse, N. Y. 13201

USDA Reg. No. 8590-59

