

THIRAM 4.8 D

NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENT:
Thiram (tetramethyl thiuram disulfide) 4.875%
INERT INGREDIENTS: 95.125%
100%

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

May cause irritation of eyes, nose, throat, and skin. May be harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin and eyes. In case of contact, wash with plenty of soap and water; for eyes, flush with water and get medical attention.

This product is toxic to fish. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

SP 100

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

APPLES: To control scab, fly speck, sooty blotch, bitter rot, Brock's spot, and cedar-apple rust. Apply 1 to 2 pounds per average size tree. To control scab, start applications in pre-pink and repeat before or early in infection (wet) periods through pink, bloom, petal-fall, and cover periods. To control cedar-apple rust, make additional applications if necessary to keep trees protected during the bloom period.

PEACHES: To control brown rot, Rhizopus rot, and scab, apply 1/2 to 1 pound per average size tree. Apply at 3 to 4-day intervals during bloom for blossom blight control. Apply at petal fall and at 10 to 14-day intervals thereafter for fruit rots and scab.

STRAWBERRIES: To control gray mold (Botrytis) fruit rot-apply 40 pounds per acre. Make 3 to 5 applications at 10-day intervals beginning at early blossom stage. Good coverage of buds, blossoms and young fruit is essential.

Do not use on peaches within 7 days of harvest. If applied to strawberries within 3 days of picking, remove residues by washing.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Code No. AC63E
USDA Reg. No. 8590-63

Manufactured by Agway Inc., Syracuse, N. Y.

