



# 7-1 D

## NET WEIGHT FIFTY POUNDS

### ACTIVE INGREDIENTS

cyfluthrin	1.0%	
Other Cube Resins	2.0%	
Copper (in Basic Copper Sulfate)		
expressed as metallic copper	7.0%	10.0%
<b>INERT INGREDIENTS</b>		
not more than		90.0%
	Total	100.0%

## CAUTION:

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. IRRITANT TO EYES AND SKIN. HARMFUL IF SWALLOWED.

This product is toxic to fish. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with soil or burning. Stay away from smoke.

**WARRANTY:** Seller warrants that the product conforms to its label description and is reasonably fit for the purpose stated on the label when used in accordance with directions or normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular use, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or in conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential or commercial damages, such as lost profits or value of crop.

## DIRECTIONS

READ THESE DIRECTIONS CAREFULLY. WARRANTY OF SALE EXTENDS ONLY TO ACCORDANCE WITH LABEL CAPTIONED DIRECTIONS AND DIRECTIONS.

Thorough thin even coverage of entire plant, including all leaf surfaces is essential for good insect and disease control. Do not apply dust in excess. See rate below.

**BEANS:** To control Mexican bean beetle, downy mildew, and rust, apply when insect or disease first threatens. Repeat at weekly intervals. During periods of cool, wet weather, produces rapidly growing succulent foliage occasional injury to foliage may result. Do not apply one day to harvest.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** To control diamond back moth, imported cabbage worm, downy mildew, apply when insect or disease first threatens. Repeat at weekly intervals. Do not apply one day to harvest.

**CELERY:** To prevent early and late blights begin application when plants emerge. Apply at 7-10 day intervals throughout season except during very dry weather. Do not apply one day to harvest.

**CUCUMBERS, MELONS AND SQUASH:** To control cucumber beetles, young squash bugs, squash vine borer, downy mildew, anthracnose, downy mildew, angular leaf spots, scab, and in the prevention of bacterial wilt, apply at weekly intervals beginning when plants emerge. Do not apply one day to harvest.

**POTATOES AND TOMATOES:** To control flea beetle, Colorado potato beetle, tomato horn worm and repel cutworms, apply when plants emerge. To control early and late blights on potatoes, foliage blights, and late blight on tomatoes, apply at weekly intervals. On potatoes start when plants emerge, and on tomatoes start after transplanting. Do not apply one day to harvest.

**RATE AND TIMING OF APPLICATION:** Apply 3 1/2 pounds per acre to give uniform coverage to the plants. Do not apply closer to harvest than one day.

Do not use this dust on corn or grain crops.

**NOTE:** This pesticide is for use only in New York State. See label for complete directions.

Code No. 7-1 D

USE ONLY FOR THE PURPOSES

Manufactured by Agway Inc., Syracuse, N. Y.