

PARATHION 2 D

NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENTS:

| | |
|---|---------------|
| Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate) | 2.00% |
| INERT INGREDIENTS, not more than | <u>98.00%</u> |
| | 100.00% |



DANGER:

KEEP OUT OF REACH OF CHILDREN.

May be Fatal if Swallowed, Inhaled, or Absorbed Through Skin! Rapidly Absorbed Through Skin! Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the USDA for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and food-stuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product

contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSEL SPROUTS, KALE, TURNIPS: To control aphids apply at 25 pounds per acre as soon as these insects make their appearance. Be sure to direct dust to the underside of the leaves. Repeat application in 7 to 10 days if necessary. Do not apply to cabbage, Kale and turnips within 10 days of harvest. Do not apply to cauliflower, broccoli, or brussel sprouts within 7 days of harvest.

ONIONS: To control thrips, apply at 15 to 20 pounds per acre as soon as these insects are seen. Repeat applications as necessary. Do not apply within 15 days of harvest.

BEANS: To control tarnished plant bug, Mexican bean beetle, leafhopper, aphids, two spotted mite. Apply at 25 to 40 pounds per acre. Do not apply within 15 days of harvest.


POTATOES: To control Aphis. Apply at 25 to 40 pounds per acre when aphids appear. Repeat applications as necessary. Do not apply within 5 days of harvest.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Code No. AC49E

USDA Reg. No. 8590-49

Manufactured by Agway Inc., Syracuse, N. Y.



CAPTAN-GUTHION 10-6 D


ACTIVE INGREDIENTS:

Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide) 10.0%

O, O-Dimethyl S-4-oxo-1, 2, 3-benzotriazin-3 (4H)-ylmethyl phosphorodithioate 6.0%

INERT INGREDIENTS: 84.0%

*U.S. Patent No. 2,758,115 100.0%



DANGER:

KEEP OUT OF REACH OF CHILDREN.

ANTIDOTE: If Swallowed: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY! If on Skin: In case of contact, remove contaminated clothing and immediately wash skin with soap and water. Poisonous if Swallowed, Inhaled, or Absorbed Through Skin! Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing, and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for guthion protection. Keep all unprotected persons out of the operating area or vicinity where there may be any danger of drift. Vacated areas should not be re-entered for at least two days. Do not store near feed or foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

NOTE TO PHYSICIAN: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by burying with wastes or burning. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

APPLES: To aid in control of scab, Brook's fruit spot, sooty blotch, fly-speck, coddling moth, plum curculio, leaf rollers, bud moth, aphids, and mites, apply at 7-day intervals or as conditions warrant during pre-blossom period and after petal fall, up to 15 days of harvest. Do not apply during blossom period to avoid injuring bees. Full coverage applications, 20 lbs. per acre by airplane or ground duster are necessary for satisfactory control. Do not apply more than 8 times per season.

CAUTION: Do not apply within 15 days of harvest. If disease infection periods occur within 15 days of harvest, apply a suitable fungicide as required. Do not apply to apples within 10 days of an oil spray. Use cautiously on Red Delicious and Baldwin varieties, inasmuch as wettable powder sprays with captan have sometimes caused foliage injury to these varieties. Highly Toxic to fish: Do not contaminate any body of water.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Agway Inc., Chemical Division
Box 1333, Syracuse, N. Y. 13201

EPA Reg. No. 8590-71

NET CONTENTS FIFTY POUNDS