

CAPTAN 5 D

NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENT:

Captan (1-(2-chlorophenyl)
thio)-4-cyclohexene-1,
2-dicarbonimide

5.0%

INERT INGREDIENTS:

95.0%

100.0%

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Do not inhale dust. Avoid contact with skin. Wear respirator while handling or applying dust. Avoid contamination of feed and food-stuffs.

This product is toxic to fish and wildlife. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

SP370

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF USE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Captan 5D is a fungicidal dust formulation suitable for use on many crops and ornamentals.

APPLES: Apple Scab, Black Rot, Fly Spots, Leaf Spot, Black Rot, Botrytis, etc. 30 to 35 lbs. per acre. Make first application as soon as scab spots are ready to discharge and an infection period begins. Repeat application for each subsequent infection period. Do not apply to apples within 10 days of an oil spray. Use cautiously on red delicious and Baldwin apples inasmuch as captan sprays have sometimes caused injury to the foliage of these varieties.

PEACH, NECTARINES: Brown Rot and Peach Scab, 50 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary. May cause slight injury on some varieties in the later sprays.

CELERY: (Field and Seed Bed Only): Pink Rot, Damp-off Disease. Use 30 to 40 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

CANTALOUPE, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS: Downy Mildew, Angular Leaf Spot, Anthracnose. Apply at 35 lbs. per acre on young plants, increasing to 50 to 60 lbs. per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 10 day intervals until danger of infection is past.

ORNAMENTALS: ROSES, Black Spot; CARNATIONS, Rust and Leaf Spot; CHRYSANTHE-

MUMS, Flower Blight. Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

POTATOES (Seed Pieces): Seed Rot and Damp-off. Dust at 1 1/2 to 100 lbs. cut seed pieces. Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging.

ROSES AND GREENHOUSE BENCH TREATMENT: Preplanting for Damp-off and Root Rot Diseases on Seedling or Transplants of Roses (and other shrubs and trees). Many flowers and vegetables (such as Beans, Peas, Cucurbits) and lawn seed-beds—1 1/2 to 2 1/4 lbs. per 1000 sq. ft. (50 to 60 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

CHERRIES: To control leaf spot and brown rot, apply 35 to 40 lbs. per acre. On sweet cherries, apply in pink, bloom, and subsequent sprays, at 10-day intervals or less. On sour cherries, begin applications at petal-fall. Under severe infection conditions, shorten the interval between applications.

STRAWBERRIES: To control Botrytis fruit rot, apply 30 to 40 lbs. per acre, beginning when new growth starts and repeat at 6 to 10-day intervals. Use shorter intervals under conditions favorable to infection. Continue applications up to 3 days before first picking and treat immediately after each picking.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

USDA Reg. No. 8590-92

Manufactured by Agway Inc., Chemical Division,
Box 1333, Syracuse, N. Y. 13201