


# AGWAY CAPTAN 5D

FOR USE IN CONTROLLING CERTAIN FRUIT, VEGETABLE AND ORNAMENTAL DISEASES.

## ACTIVE INGREDIENT

Captan (N-trichloromethyl) thio-4 cyclohexene-1, 2-dicarboximide	4.73%
Related Compounds	0.27%
<b>INERT INGREDIENTS</b>	<b>95.00%</b>
<b>TOTAL</b>	<b>100.00%</b>



KEEP OUT OF REACH OF CHILDREN.

## CAUTION

See additional precautionary statements below.

## DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONS AND DIRECTIONS

Captan 5D is a fungicidal dust formulation suitable for use on many crops and ornamentals.

**APPLES:** Apple Scab, Brooks Fruit Spot, Fly Specking, Bitter Rot, Black Rot, Botryosphaeria. 30 to 35 lbs. per acre. Make first application as soon as scab spores are ready to discharge and an infection period threatens. Repeat application for each subsequent infection period. Do not apply to apples within 10 days of an oil spray. Use cautiously on Red Delicious and Baldwin apples inasmuch as Captan sprays have sometimes caused injury to the foliage of these varieties.

**PEACH, NECTARINES:** Brown Rot and Peach Scab 50 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary. May cause slight injury on some varieties in the later sprays.

**CELERY:** (Field and Seed Bed Only): Pink Rot, Damp-off Disease. Use 30-40 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

**CANTALOUPE, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS:** Downy Mildew, Angular Leaf Spot, Anthracnose. Apply at 35 lbs. per acre on young plants, increasing to 50-60 lbs. per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 10 day intervals until danger of infection is past.

**ORNAMENTALS:** ROSES, Black Spot; CARNATIONS, Rust and Leaf Spots; CHRYSANTHEMUMS, Flower Blight. Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

**POTATOES (Seed Pieces):** Seed Rot and Damp-off. Dust at 1½ lb. to 100 lbs. cut seed pieces. Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days, the treated pieces should be stored for 2 days in open crates before bagging. Do not use treated seed pieces for food or feed.

**SOIL AND GREENHOUSE BENCH TREATMENT:** Preplanting for Damp-off and Root Rot Diseases on Seedling or Transplants of Roses (and other shrubs and trees), many flowers and vegetables (such as Beans,

Peas, Cucurbits) and lawn seedbeds — 1½ to 2¼ lbs. per 1000 sq. ft. (50 to 60 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

**CHERRIES:** To control leaf spot and brown rot, apply 35-40 lbs. per acre. On sweet cherries, apply in pink, bloom, and subsequent sprays, at 10 day intervals or less. On sour cherries, begin applications at petal-fall. Under severe infection conditions, shorten the interval between applications.

**STRAWBERRIES:** To control Botrytis fruit rot, apply 30-40 lbs. per acre, beginning when new growth starts and repeat at 6 to 10 day intervals. Use shorter intervals under conditions favorable to infection. Continue applications up to 3 days before first picking and treat immediately after each picking.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

**CAUTION:** Causes skin and eye irritation. May be harmful if swallowed or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Use with adequate ventilation. Wash thoroughly after handling.

**FIRST AID:** In case of eye contact, immediately flush eyes with plenty of water for at least 15 minutes. Call a physician. In case of skin contact, wash with soap and water.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

**DISCLAIMER:** Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

**NOTE:** This pesticide is to be sold ONLY in this original unbroken container.

EPA Reg. No. 8590-92

EPA Est. No. 8590-PA-1

AGWAY INC., CHEMICAL DIVISION  
BOX 4933, SYRACUSE, NY 13221

Net Contents 50 lbs.

A 7 81