

ACCEPTED

OCT 13 1967

CHLORDANE 5D

Net Weight 50 lbs.

ACTIVE INGREDIENT:

*Technical Chlordane

INERT INGREDIENTS: Not more than

5.0

95.0

100.0

*Equivalent to 3,8-octachloro-4,7-methanotetrahydroindane and 2 isomers of related compounds.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

CAUTION KEEP OUT OF REACH OF CHILDREN.

Avoid inhalation. Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Do not contaminate any body of water or use for purposes other than those shown on the label.

STATEMENT OF LIABILITY: Agway warrants that this material conforms to the chemical description on the label and is fit for the purposes referred to in the directions, subject to the inherent risks referred to therein. Agway makes no other express or implied warranty of fitness or of merchantability, and no agent of Agway is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to the purchase price and shall not include consequential damages.

DIRECTIONS

Because of the range of conditions under which this product may be used, it is impossible to eliminate all risks inherently associated with its use, such as the risk of ineffectiveness, of plant and animal injury, or of excessive residues, etc., even though the product is used in accordance with these directions. Agway offers this product subject to the understanding that the buyer assumes all such risks.

Thorough coverage of area treated is essential for adequate pest control.

LAWNS OR TURF: To control subsoil insects such as grubs of Japanese beetle, Oriental beetle, Asiatic garden beetle, masked chafer, European chafer, May beetle and rose chafer, use 5 lbs. per 1000 sq. ft. (200 lbs. per acre). To control above ground pests such as ants, chinch bugs, cutworms, earwigs, sod webworms, and sowbugs, use 2-1/2 lbs. per 1000 sq. ft. (110 lbs. per acre).

Do not treat new lawns until sod is well established to avoid injury to the seedling grasses. Apply in early spring to established turf. Do not feed treated turf to Dairy Animals or Animals being finished for slaughter.

TO AVOID INJURY TO GRASS, WATER INTO GROUND AFTER APPLICATION.

Keep children and pets off the treated area until the insecticide has been washed into the soil and the treated area has dried completely.

FLOWERS AND ORNAMENTALS: To control tansy or black vine weevil, grasshoppers, thrips and plant bugs, dust plants lightly but uniformly. Apply when insects appear and repeat as necessary. Also treat soil for 2 or 3 inches at base of plant to aid in controlling tansy weevil.

To control wireworms, cutworms, sowbugs, and grubs of Japanese beetle, Oriental beetle, Asiatic garden beetle, masked chafer, European chafer, May beetle and rose beetle, use 5 lbs. per 1000 sq. ft. (200 lbs. per acre). Use as a ground application in early spring. Where possible mix into soil by discing immediately to a depth of 4 to 5 inches.

VEGETABLE GARDEN SOIL FOR PLANTING OF BEANS, BEETS, BROCCOLI, CABBAGE, CAULIFLOWER, CHERRY, EGGPLANT, LETTUCE, BULB ONIONS, PEPPERS, POTATOES, RADISHES, TOMATOES, TURNIPS: To control cutworms, root maggots, earwigs, mole crickets, sowbugs, millipeds, ants, wireworms and grubs of Japanese beetle, Oriental beetle, Asiatic garden beetle, masked chafer, European chafer, May beetle and rose chafer, use 5 lbs. per 1000 sq. ft. (200 lbs. per acre). Apply to ground prior to planting. Mix into soil in garden beds to a depth of 4 to 5 inches. Use as a pre-planting soil application only. Do not use on ground in spring garden.

TIMING OF APPLICATION: Timing and rate of application vary with conditions. For best results, apply to ground in early spring, before the insects appear, and repeat at intervals of 3 to 4 weeks. For control of insects that appear later in the season, apply when they first appear. Do not use on ground in spring garden.

AGWAY CHEMICALS COMPANY, INC., NEW YORK, N.Y.