

# PHYGON® 3 D

NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENTS:

\*Dichlone (2,3-dichloro-1,4-naphthoquinone) ..... 3.0% by wt.  
INERT INGREDIENTS, not more than ..... 97.0% by wt.  
Total ..... 100.0%

\*Dichlone (2,3-dichloro-1,4-naphthoquinone) is otherwise known as Phygion which is a trademark of U.S. Rubber Co. and is covered by U.S. Patent No. 2,349,772.

## CAUTION:

KEEP OUT OF REACH OF CHILDREN.

May cause skin irritation. Avoid prolonged repeated contact. Avoid inhalation of dust. Use with adequate ventilation. Wear protective clothing. Wash hands and face thoroughly with soap and water after use or before eating or smoking. Keep away from feed and foodstuffs. See other warnings elsewhere on label. Toxic to fish.

This product is toxic to fish. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

## DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

This dust is especially formulated for use with air dusting or ground dusting equipment. Thorough and uniform coverage of all areas is essential for best disease control.

**APPLES:** To aid in controlling apple scab during the primary scab period, use 30 to 40 lbs. per acre. Apply prior to infection periods to prevent scab and up to 48 hours after an infection period begins (depending on temperatures) to control scab infections. Repeat application may be necessary following heavy rainfall. Repeated applications or applications during low temperatures may increase fruit russeting. Do not apply to apples within 1 day of harvest.

**GRAPES AND PEACHES:** To aid in controlling brown rot blossom blight, use 30 to 40 lbs. per acre during early bloom through petal-fall stages. Apply prior to infection periods to prevent brown rot blossom blight and up to 12 hours after an infection period begins to control infections. Do not apply later than petal-fall.

Timing and rate of application vary as a result of local conditions. Since these conditions are unknown to the manufacturer, you must satisfy yourself as to local requirements. Do not exceed maximum rates or apply later than directed above.

**WARNING:** Do not use in combination with organic mercury fungicides or apply during high temperatures.

**NOTE:** This pesticide is to be sold ONLY in this original unbroken package.

Code No. AC122E

USDA Reg. No. 8590-122

Manufactured by Agway Inc., Syracuse, N.Y.

