

MITICIDE 2 D

Contains 2% Morestan*


ACTIVE INGREDIENT:	
6-Methyl-2, 3-quinoxalinedithiol cyclic S, S-dithiocarbonate**	2.0%
INERT INGREDIENTS:	98.0%
-	
1	00.0%

*Reg. U.S. Pat. Off. by Farbenfabriken Bayer, A. G., Chemagro Corporation licensee.

**U.5. Patent No. 3,091,613.

-





KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. May cause skin irritation. Contact with skin, eyes, or clothing should be avoided. Avoid breathing of dust. Wash thoroughly with soap and warm water after handling. Do not contaminate feed or foodstuffs.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. A : 'y this product only as specified on this label.

Do not reuse empty container. Destroy by burying with wastes or burning. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS **READ THESE ENTIRE DIRECTIONS AND WARRANTY** OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

APPLES, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES, PEARS: Pre-Bloom for control of mites and mite eggs. Apply 30 to 50 lbs. per acre in pre-bloom period (delayed dormant, pre-pink or pink). Do not apply after first bloom.

APPLES, PEARS: Post-Bloom for control of mites and mite eggs. Apply 30 to 50 lbs. per acre. Repeat as necessary but NOT MORE THAN 3 TIMES PER SEASON. Do not apply within 35 days of harvest. Foliage and fruit injury may occur in some locations, under certain conditions (particularly Anjaw variety from summer applications). Do not graze treated areas. NOTE: This pesticide is to be sold ONLY in the original unbroken pack-

Agway Inc., Chemical Division Box 1333, Syracuse, N. Y. 13201

age.

