

Net Contents



50 Pounds

CAPTAN-GUTHION-LEAD 5-1.5-7 D

(For Control of Certain Insects and Diseases on Apples)

ACTIVE INGREDIENTS:

Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	5.00%
*O,O-Dimethyl S-[4-oxo-1,2,3-benzotriazin-3(4H)ylmethyl] phosphorodithioate	1.50%
Lead Arsenate	7.00%
1.40% Total Arsenic (As Metallic)	
0.14% Water-Soluble Arsenic (As Metallic)	

INERT INGREDIENTS: 86.50%

*U. S. Pat. No. 2,758,115

WARNING: KEEP OUT OF REACH OF CHILDREN.

May Be Fatal If Swallowed, Inhaled, or Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Avoid breathing dust or spray mist. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type approved by U. S. Bureau of Mines for guthion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be reentered for at least two days. Do not store near feed and food products. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke or fumes.

Not for use or storage in or around the home.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

A fungicide-insecticide mixture for controlling certain diseases and insect pests affecting apples.

APPLES: Apply Scab, Brooks Fruit Spot, Fly Speckling Bitter Rot, Black Rot, Bud Moth, Codling Moth, Plum Curculio, Red-banded Leaf Roller, Fruit Tree Leaf Roller, Eye-spotted Bud Moth, Mites (European Red Mite, Apple Rust Mite, Two-spotted Mites) and Aphids (Rosy Apple Aphid, Green Apple Aphid, Woolly Apple Aphid).

Apply during pre-bloom period and at petal fall at 7 to 10 day intervals or as necessary for disease and insect control.

Continue applications during the growing season at 10 to 14 day intervals as necessary for disease and insect control. Special sprays or dust applications may be necessary to control mites.

Rate of Application: Apply 35 to 40 lbs. per acre or 1 to 2 lbs. per mature apple tree. Full-coverage applications using power sprayers are necessary for satisfactory control.

WARNING:

Do not apply during bloom period to avoid injury to bees.

Do not apply to apples within 10 days of an oil spray.

Do not apply after the 5th cover or within 15 days of harvest on apples.

Use cautiously on Red Delicious and Baldwin apples during early application inasmuch as Captan may cause injury to the foliage of these varieties.


Remove excess residues at harvest.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Manufactured by Agway Inc. - Chemical Division, Box 1333, Syracuse, New York 13201

8590-132



USDA Reg. No. 8590-132