

CAPTAN 10 D

OCT 3, 1967

Not for use on...

AGWAY INCORPORATED  
1000 N. 10th St., P.O. Box 1000, St. Paul, Minn. 55101

USE CAPTAN 10 D ONLY IN ACCORDANCE WITH LABEL DIRECTIONS, WARNINGS AND PRECAUTIONS.

**CAUTION KEEP OUT OF REACH OF CHILDREN.**

Avoid prolonged or repeated contact with skin. Wear a respirator while handling or applying dust. Avoid contamination of feet and foodstuffs. Rinse spray equipment and containers and dispose of liquid wastes in a proper manner. Do not allow to flow away from water supplies.

Keep out of any body of water. Fish may be killed.

**STATEMENT OF LIABILITY:** Agway warrants that this material conforms to the chemical description on the label and is fit for the purposes referred to in the directions subject to the inherent risks referred to therein. Agway makes no other express or implied warranty of fitness or of merchantability, and no agent of Agway is authorized to do so except in writing, with specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to the purchase price and shall not include consequential damages.

**DIRECTIONS**

Because of the range of conditions under which this product may be used, it is impossible to eliminate all risks inherent in its use, such as, the risk of ineffectiveness, of plant and animal injury, or of excessive resistance to it, even though the product is used in accordance with these directions. Agway offers this product subject to the understanding that the buyer assumes all such risks.

Captan 10 D is a fungicidal dust formulation suitable for use on many crops and ornamentals.

**APPLES:** Apple Scab, Brocks Fruit Spot, Fly Specking, Bitter Rot, Black Rot, Botryosphacteria, Fire Blight, Pear Scab. Make first application as soon as scab spores are ready to discharge and an infection period threatens. Repeat applications at 10 to 14 day intervals. Do not apply to apples within 10 days of an apple spray. Use caution on Baldwin apples inasmuch as Captan sprays have sometimes caused injury to the foliage of these varieties.

**PEACH, NECTARINES:** Brown Rot and Peach Scab. 25 lbs. per acre. Make 2 to 3 pre-harvest applications. Repeat at intervals of 10 to 14 days as necessary. Make pre-harvest application when blossoms are in full bloom. Make final application 10 to 14 days before harvest.

**CHERRY:** Brown Rot, Cherry Leaf Spot, Dampest Disease. Use 40 to 50 lbs. per acre or 1 lb. per 1000 sq. ft. Apply when blossoms are in full bloom. Repeat as necessary.

**CANTALOUPES, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS:** Downy Mildew, Angular Leaf Spot, Anthracnose. Apply at 20 lbs. per acre on young plants, increasing to 25 to 30 lbs. per acre on mature plants. Apply at first sign of disease. Repeat at 10 to 14 day intervals until danger of infection is past.

**CANNA LILIES, PEONIES, IRIS, SPINACH, CARNATIONS, RIBS and Leaf Spot, CHRYSANTHEMUMS, Flower Fungus.** Apply at 20 lbs. per acre or 1 lb. per 1000 sq. ft. at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 10 to 14 day intervals as necessary.

**FRUIT TREES:** Brown Fruit Rot, Apple and Leaf Spot, and Root Rot. Apply at 2 to 4 lbs. per 1000 sq. ft. Repeat at 10 to 14 day intervals as necessary. Use Captan 10 D on fruit trees in the spring. Make first application at first sign of disease. Repeat at 10 to 14 day intervals as long as possible with weather permissable.

**POTATOES (See Peaches):** See Rot and Dampest Disease. Apply at 20 to 30 lbs. per acre. Repeat at 10 to 14 day intervals. Do not apply to potatoes until 2 weeks before harvest. Do not apply to potatoes within 10 days of harvest. Do not apply to potatoes within 2 weeks of the time the tubers are to be used for seed.

**SCALES AND GREENHOUSE BENCH TREEMENDS:** Apply at 20 to 30 lbs. per acre. Repeat at 10 to 14 day intervals. See label on the spray for directions on other fruits and trees. Note: Captan 10 D is not for use on plants in greenhouses. Do not apply to plants in greenhouses. Do not apply to plants in greenhouses.

**CHRYSANTHEMUMS:** Apply at 20 to 30 lbs. per acre. Repeat at 10 to 14 day intervals. Do not apply to chrysanthemums within 10 days of harvest. Do not apply to chrysanthemums within 10 days of harvest. Do not apply to chrysanthemums within 10 days of harvest.

**STRAWBERRY:** Apply at 20 to 30 lbs. per acre. Repeat at 10 to 14 day intervals. Do not apply to strawberries within 10 days of harvest. Do not apply to strawberries within 10 days of harvest. Do not apply to strawberries within 10 days of harvest.

For more information, contact your local Agway office.