

ontents: 6 Ounces

Makes one gallon of liquid bait,

DUOCIDE Soluble Bait

Kills Rats and Mice

CTIVE INGREDIENTS:

Sodium salt of 2-Pivalyl-1, 3-Indandione*

ERT INGREDIENTS:

-Equivalent to 2-Pivalyl-1, 3-Indandione 0.13%

J. S. Patent No. Re. 23,080.

0, 14% 99, 86%

CAUTION KEEP OUT OF REACH OF CHILDREN.

he active ingredient in Duocide Soluble Bait is pivalyn, an anti-coagulant chemical. Baits prepared from it must protected or placed where humans, livestock, poultry, wildlife, or pets will not eat or drink them. If swallowed humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding, acase of eating or drinking, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is lear. Call a physician immediately.

OTE TO PHYSICIAN: Administration of blood transfusions combined with intravenous injections and oral doses of itami. 1 is indicated.

'ARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose ated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor by other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this oduct contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to ller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited direct damages, and shall no include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE FITTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL AUTIONS, WARNINGS AND DIRECTIONS.

o prepare liquid bait, simply dissolve the contents of this package in 1 gal. of clean water. Pour liquid bait into ass, plastic, or crockery dish or chicken fount. (Metal containers are not recommended.) Prepare only the amount liquid bait which is intended for exposure at any one time.

acement of Baits - Place baits along walls, in corners, and in concealed locations where rats and mice can feed undisturbed. Baits should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes. Continue baiting until all feeding has stopped. Immediately replace liquid baits that become cloudy.

- Pint liquid dispenser per location should be the minimum amount placed. An uninterrupted supply of bait should be maintained for at least 10 days.

- Pint liquid dispenser should be placed at 8 to 10 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

OTE: This pesticide is to be sold ONLY in this original unopened package.

ANUFACTURED FOR AGWAY INC. - CHEMICAL DIVISION, BOX 1333, SYRACUSE, NEW YORK 13201

ACCEPTED

ats

ice

JUL 8 1971



USDA Reg. No. 8590-148