श्विकृत

FOR: AGWAY MANEB-THIODAN 4.5-3D **DATE:** 11/2/82 (((FRONT PANEL))) NET WT. PAR SCHAPAUTERS -- UTE = JUSTAPAUTER 1#, 50# **AGWAY** MANEB-THIODAN 4.5-3D To control insects & diseases of Vegetables ACTIVE INGREDIENT: Maneb (manganese ethylene bisdithiocarbamate) 4.5% (metallic manganese content 0.98%) Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide) 3.0% INERT INGREDIENTS: 92.5% TOTAL 100.0% KEEP OUT OF REACH OF CHILDREN CAUTION (18 pt.) See additional precautionary statements on side panel. O'K BY AGWAY INC. CHEMICAL DIVISION BOX 4933 EPA Reg. No. 8590-164 SYRACUSE, NY 13221 EPA Est. No. 8590-PA-1

SHEET NO.1 - 8

ı

4

GOBA

FOR:

4.1

AGWAY MANEB-THIODAN 4.5-3D

DATE: 11-2-82

PRECAUTIONARY STATEMENTS

71CA 0.0 2 + 4442 184 , to 18 + 40 C 444 Au 144

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Do not breath dust. Avoid contact with skin, eyes or clothing. In case of contact with eyes, flush with plenty of water. Get medical attention.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air. Apply artificial respiration, if indicated.

If on skin, remove contaminated clothing & wash affected areas with soap & water.

If in eyes, flush eyes with plenty of water. Call a physician if irritation persists.

Wash hands thoroughly after using & before eating or smoking. During prolonged or commercial exposure, wear a pesticide respirator jointly approved by the Mining Enforcement & Safety

Administration (formerly US Bureau of Mines) and by the National Institute for Occupational Safety & Health under provisions of 30 CFR Part 11.

O'K BY

SHEET NO. 2 - 8

Radio S

FOR:

30

AGWAY MANEB-THIODAN 4.5-3D

DATE: 11/2/82

Pica giranacteni Fuiteriochanacteni

Wear protective clothing (long pants, long sleeve shirt, gloves, hat & boots) during mixing & loading.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds & other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated crop. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

O.K BA

STORAGE & DISPOSAL (12 pt.)

1# Do not reuse empty container. Wrap container
and place in trash collection.

SHEET NO. 3 - 8

FOR: AGWAY MANEB-THIODAN 4.5-3D **DATE:** 11/2/82

Do not contaminate water, food or feed by storage or disposal.

50# Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation & Recovery Act.

Completely empty bag by shaking & tapping sides & bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bag in a sanitary landfill or by incineration if allowed by state & local authorities.

This product is formulated to control both insects and diseases. Best results can be obtained by applying before serious build-up occurs. Repeat at 7 to 10 day intervals throughout the growing season. Make thorough coverage of top and underside of foliage. Number of days between treatment & harvest are indicated after crop in ().

BEANS (snap, string, lima, dry) (4) - For control of Anthracnose, downy mildew, rust, cucumber beetles, leafhoppers, Mexican bean beetles, green stink bugs, black bean aphids & bean leaf skeletonizers, apply 30 lbs. dust per acre (1/2# per 725 sq. ft.). Do not apply to lima beans that

SHEET NO. 4 - 8

<u>eopy</u>

FOR:

AGWAY MANEB-THIDAN 4.5-3D

DATE: 11/2/82

Pak jaguahuste

are to be removed from field for processing. Do not feed treated threshings to livestock or allow to graze in treated fields. Do not apply more than 3 times per season.

BROCCOLI (7), BRUSSELS SPROUTS (14), CABBAGE (7), CAULIFLOWER (14) - For control of Alternaria leaf spot, downy mildew, aphids, cabbage loopers, imported cabbageworms, Diamond-back moths, larvae, cross-striped cabbageworms, flea beetle, Harlequin bugs and stink bugs, apply 30 lbs. dust per acre (1/2# per 725 sq. ft.).

CARROTS (7) - For control of Alternaria leaf spot, cercospora leaf spot, flea beetles, leaf-hoppers, lygus bugs, spittlebugs and stink bugs, apply 20 to 40 lbs. dust per acre (1/2 to 1 lb. per 2,200 sq. ft.). Do not apply more than twice per season. Do not use tops for food or feed.

O'K BY

CUCUMBERS (5), MELONS (5), including Cantaloupes, Cassabas, crenshaws, honeydews, muskmelons, Persian melons and watermelons, SQUASH (5), Summer and winter, PUMPKINS (0) - For control of Alternaria leaf spot, angular leaf spot, anthracnose, cercospora leaf spot, downy mildew, gummy

SHEET NO.5 - 8

<u>aapy</u>

FOR:

1

AGWAY MANEB-THIODAN 4.5-3D

DATE: 11/2/82

PART OF LANGUAGES AND THE

stem blight, Pythium fruit rot, scab, aphids, cucumber beetles, melonworms, pickleworms, squash beetles, squash bugs, squash vine borers, and striped cabbage flea beetles, apply 30 lbs. dust per acre (1/2# per 725 sq. ft.).

LETTUCE (14) including head & leaf - For control of downy mildew, aphids, cabbage loopers, Imported cabbageworms, leafhoppers, & tarnished plant bugs, apply 30 lbs. dust per acre (1/2# per 725 sq. ft.). For head lettuce, do not apply more than 3 times after thinning; remove outer leaves at harvest. For leaf lettuce, do not make more than 2 applications per season.

POTATOES (0) - For control of early & late blight, aphids, Colorado potato beetles, flea beetles & leafhoppers, apply 33 lbs. dust per acre (1/2 lb. per 660 sq. ft.). Do not plant root crops other than carrots, potatoes or sweet potatoes as follow up crops. As plants become larger & late blight threatens, apply 40 lbs. dust per acre (1/2# per 550 sq. ft.). Apply at 7 day intervals. If late blight is present, shorten interval to 3 to 5 days.

0'K 8Y

SHEET NO. 6 - 8

FOR: AGWAY MANEB-THIODAN 4.5-3D

DATE: 11/2/82

SPINACH (21) - For control of downy mildew, aphids a spinach leaf miners, apply 25 lbs. dust per acre (1/2* per 875 sq. ft.). One application per season.

SWEET CORN (0) - For fresh vegetable use only.

For control of Helminthosporium leaf blight, Corn
leaf aphids and corn earworms, apply 50 lbs. dust
per acre (1/2# per 440 sq. ft.). Do not apply to
sweet corn to be processed. Do not make more than
5 applications per season. Do not feed forage or
ensilage to livestock or allow livestock to
graze in treated fields.

TOMATOES (1), EGGPLANTS (1), PEPPERS (1) - For control of anthracnose, cercospora leaf spot, early & late blight, gray leaf spot (Stemphylium, Septoria leaf spot, aphids, blister beetles, Colorado potato beetles, flea beetles, green stink bugs, tomato fruitworms, tomato hornworms and whiteflies, Apply 30 lbs. dust per acre (1/2# per 725 sq. ft.). As plants become large & late blight threatens, apply 40 lbs. dust per acre (1/2# per 550 sq. ft.). Apply at 7 day intervals. If late blight is present, shorten interval to 3 to 5 days.

O'K BY

SHEET NO. 7 - 8

FOR: AGWAY MANEB-THIODAN 4.5-3D DATE: 11/2/82

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

РОА ОСНЦЯАЦІТЕЯ: ПОТЕН ФОНЦЯАСТЕВН

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

O'K BY

SHEET NO. 8 - 8