

DODINE 1 D

Contains Cyprex* for Control of Certain Diseases in Apples,
Roses and Strawberries

ACTIVE INGREDIENTS:

Dodine (n-dodecylguanidine acetate) 1.00%

INERT INGREDIENTS: 99.00%

100.00%

*TM of American Cyanamid Co., Reg. U.S. Pat. No. 24142

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Concentrated material causes eye irritation. In case of contact with eyes, flush with plenty of water for at least 15 minutes; get medical attention. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not re-use empty container. Destroy it by burying with wastes or burning. Stay away from smoke or fumes.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

SP 477

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

For best results, apply when air is calm, covering foliage with a light film of dust. Apply as a preventive program making the first application before disease appears. Repeat applications as necessary. Keep new growth protected. This product is suitable for application through ground dusters, airplane or hand equipment.

APPLES: Including Flower Crab Apples. For control of Apple Scab. Apply 60 to 80 lbs. of dust per acre or approximately 2 to 2½ lbs. of dust per mature apple tree. Make the first application during pre-bloom period. Repeat at 5 to 7 day intervals through first cover. Do not apply within 5 days of harvest. Do not graze cover crops in treated orchards. Do not use treated apples in the manufacture of apple pomace for use in livestock feeds.

ROSES: For control of Black Spot. Apply 30 to 40 lbs. per acre or ¾ to 1 lb. per 1,000 sq. ft. Make the first application when young leaves first appear. Repeat at 7 to 10 day intervals throughout the season. Thorough coverage as necessary. Dust lightly, do not over-treat. Leaf injury has been reported on certain rose varieties and at excessive rates.

STRAWBERRIES: For control of Leaf Scorch, Leaf Spot and Leaf Blight. Apply 90 to 100 lbs. per acre or 2 to 2½ lbs. per 1,000 sq. ft. Make the first application when blossoms first appear and repeat at 7 day intervals as necessary. Do not apply within 14 days of harvest. Do not apply in combination with lime or chlorobenzilate. Avoid drift of dust to adjacent crops. Do not graze cover crops in treated areas. Consult your State Agricultural Experiment Station or State Extension Service Specialist for proper timing of applications.

NOTE: This pesticide is to be used ONLY in this original unbroken package.

Agway Inc., Chemical Division
Box 1333, Syracuse, N. Y. 13201

EPA Reg. No. 8590-165

NET CONTENTS FIFTY POUNDS