




COPPER METHOXYCHLOR 7-5 D

For Control of Certain Diseases and Insects on Grapes

ACTIVE INGREDIENTS:

Copper (in Basic Copper Sulfate) expressed as metallic copper	7.00%
*Methoxychlor	5.00%
INERT INGREDIENTS:	88.00%
	<hr/>
	100.00%

*Equivalent to 4.4% 2, 2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 0.6% other isomers and related compounds.



CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Avoid breathing dust.

Highly toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift or runoff from the crop area treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke or fumes.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions

of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

A fungicide, insecticide dust mixture formulated for use by aerial application on Grapes. Complete coverage of all leaves including new growth, is essential for satisfactory control of Black Rot, Downy Mildew, Powdery Mildew, Grape Berry Moth, Grape Leafhopper and Japanese Beetle. Apply 30 to 40 lbs. per acre. Make first application at petal fall and repeat in 7 to 14 days. For second brood of Berry Moth, apply 35 to 45 days after bloom or when eggs are found repeat in 14 days. For diseases apply when conditions are favorable for development and repeat at 7 to 10 day intervals as necessary for control. Timing and rate of application vary according to local conditions. Consult State Agricultural Extension Service for timing and rates. Do not apply to grapes within 14 days of harvest.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

Agway Inc., Chemical Division
Box 1333, Syracuse, N. Y. 13201

EPA Reg. No. 8590-166

NET CONTENTS FIFTY POUNDS