

COPPER - PARATHION 7-2D

Net Weight 50 lbs.

For Control of Certain Diseases and Insects on Grapes


ACTIVE INGREDIENTS:

Copper (in Basic Copper Sulfate) expressed as metallic copper ----- 7.0%

Parathion (O, O-diethyl O-p-nitrophenyl thiophosphate) ----- 2.0%

INERT INGREDIENTS: ----- 91.0%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

**POISON****DANGER KEEP OUT OF REACH OF CHILDREN.**

May be fatal if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the USDA for Parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticides and volatile residues have dissipated. Do not contaminate feed and food-stuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke.)

STATEMENT OF LIABILITY: Agway warrants that this material conforms to the chemical description on the label and is fit for the purposes referred to in the directions, subject to the inherent risks referred to therein. Agway makes no other express or implied warranty of fitness or of merchantability, and no agent of Agway is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to the purchase price and shall not include consequential damages.

DIRECTIONS

Because of the range of conditions under which this product may be used, it is impossible to eliminate all risks inherently associated with its use, such as, the risk of ineffectiveness, of plant and animal injury, or of excessive residues, etc., even though the product is used in accordance with these directions. Agway offers this product subject to the understanding that the buyer assumes all such risks.

A fungicide, insecticide dust mixture formulated for use by aerial applicators on grapes. This dust contains Parathion which is not stable as a dust and should be used within one growing season.

GRAPES: For control of Black Rot, Grape Berry Moth, Mealybug, Grapevine Aphid, Japanese Beetle, Red-Banded Leaf Roller and Two-Spotted Spider Mite. Apply 25 to 30 lbs. per acre. For Black Rot control make the first application pre-bloom just before blossoms open. Repeat as a post-bloom application immediately after bloom and in 7 to 10 days after post-bloom application. For insect control, apply when insects are noticed and repeat as needed. Do not apply within 14 days of harvest.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Manufactured by AGWAY Inc. Syracuse, New York
AC-172-F

USDA Reg. No. 8590-172

APR 27 1967