

ACCEPTED

DEC 21 1984

Under Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 8590-174

AGWAY

MANEB-THIODAN 5.6-3D

ACTIVE INGREDIENTS

Maneb (Manganous ethylene bisdithiocarbamate)	5.6%
Endosulfan (Hexachlorohexahydromethane-2,4-benzodioxathiepin-3-Oxide)	3.0%

INERT INGREDIENTS	91.4%
-------------------------	-------

TOTAL	100.0%
-------	--------

Thiodan is the registered Trademark of Canada Hoechst LTD.

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements on side panel.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

CABBAGE, CAULIFLOWER: For effective disease control, dusting should start when plants are small and be applied at 7 to 10 day intervals. Do not exceed the dosage or apply beyond the date indicated under each crop as listed. To control Cabbage Looper, Imported Cabbage Worm, Thrips and Downy Mildew, apply at the rate of 20 to 25 lbs. per acre. Do not apply to Cauliflower within 14 days of harvest and to Cabbage within 7 days of harvest.

BEANS: To control Mexican Bean Beetle, Downy Mildew, apply at rate of 25 to 30 lbs. per acre. Begin applications during early bloom or when disease first appears. Do not apply more than 3 times per season. Do not apply within 3 days of harvest. Do not apply to Lima Beans that are to be removed from the field for processing. Do not feed treated threshings to livestock or allow to graze in treated fields. Do not apply to Succulent Beans within 4 days of harvest.

CUCUMBERS, MELONS, SQUASH (Summer and Winter): To control Cucumber Beetle, Two-Spotted Mite, Scab, Alternaria Leaf Spot, apply 25 to 30 lbs. per acre. Do not feed crop waste to livestock. Do not apply within 5 days of harvest.

POTATOES: To control Colorado Potato Beetle, Flea Beetle, Leaf Hopper, Early Blight, Late Blight, apply at the rate of 33 lbs. per acre. Apply as soon as plants are 2 to 6 inches high. Do not plant root crops other than carrots, potatoes, and sweet corn as follow-up crops.

TOMATOES: Flea Beetles, Aphids, Colorado Potato Beetle, Hornworms, Fruit Worms, Late Blight, Early Blight, apply at the rate of 35 lbs. per acre. Apply when fruit of first clusters is well formed. Do not apply within 5 days of harvest.

DISCLAIMER: AGWAY DISCLAIMS ALL IMPLIED WARRANTIES, INCLUDING IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL INJURY, LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS FOR USE AND PRECAUTIONARY STATEMENTS. ANY DAMAGES SHALL BE LIMITED TO DIRECT DAMAGES, NOT TO EXCEED THE PURCHASE PRICE OF THE PRODUCT AND SHALL NOT INCLUDE ANY INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES.

NOTE: This pesticide is to be sold ONLY in this original unbroken package

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation. Avoid contact with eyes, skin and clothing. Avoid contamination of feed and foodstuffs. Wash with soap and water after handling and before eating or smoking; wear clean clothing.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin: Remove contaminated clothing and wash affected areas with soap and water.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Wear protective clothing (long pants, long sleeve shirts, gloves, hat and boots) while mixing and loading.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, cleaning of equipment or disposal. Open dumping is prohibited.

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures under Subtitle C of the Resource Conservation and Recovery Act.

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

EPA Reg. No. 8590-174

EPA Est. No. 8590 PA 1



AGWAY INC. — CHEMICAL DIVISION
PO BOX 4933
SYRACUSE, NY 13221

NET WEIGHT 50#