

PHYGON - FERBAM 1-3, 8D

Net Contents 50 lbs.

For Use on Apples, Cherries and Peaches for the Control of Certain Diseases
(Contains Phygion and Ferbam)

ACTIVE INGREDIENTS:

*Dichlone (2,3-dichloro-1, 4-naphthoquinone) ----- 1.0%
Ferbam (ferric dimethyl dithiocarbamate) ----- 3.8%

INERT INGREDIENTS: Not more than ----- 95.2%

*Dichlone (2,3-dichloro-1, 4-naphthoquinone) Phygion is a trademark of U. S. Rubber Co., and is covered by U. S. Patent No. 2,349,772.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

CAUTION KEEP OUT OF REACH OF CHILDREN.

May cause skin irritation. Avoid prolonged repeated contact. Avoid inhalation of dust. Use with adequate ventilation. Wear protective clothing. Wash hands and face thoroughly with soap and water after use or before eating or smoking. Keep away from feed and foodstuffs. See additional WARNING below.

This product is toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift from crop area treated.

STATEMENT OF LIABILITY: Agway warrants that this material conforms to the chemical description on the label and is fit for the purposes referred to in the directions, subject to the inherent risks referred to therein. Agway makes no other express or implied warranty of fitness or of merchantability, and no agent of Agway is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to the purchase price and shall not include consequential damages.

DIRECTIONS

Because of the range of conditions under which this product may be used, it is impossible to eliminate all risks inherently associated with its use, such as, the risk of ineffectiveness, of plant and animal injury, or of excessive residues, etc., even though the product is used in accordance with these directions. Agway offers this product subject to the understanding that the buyer assumes all such risks.

This dust is especially formulated for use with air dusting or ground dusting equipment. Thorough and uniform coverage of all trees is essential for best disease control.

APPLES: To aid in controlling apple scab as well as Apple Cedar Rust, Blossom Blight during the primary scab period, use 30 to 40 lbs. per acre. Apply prior to infection periods to prevent scab and up to 48 hours after an infection period begins (depending on temperature) to control scab infections. Repeat application may be necessary following heavy rainfall. Repeated applications or applications during low temperatures may increase fruit russetting. Do not apply within 7 days of harvest.

CHERRIES AND PEACHES: To aid in controlling brown rot blossom blight, use 30 to 40 lbs. per acre during early bloom through petal-fall stages. Apply prior to infection periods to prevent brown rot blossom blight and up to 12 hours after an infection period begins to control infections. On Cherries, do not apply within 3 days of harvest. On Peaches, do not apply within 21 days of harvest.


TIMING OF APPLICATION: Timing and rate of application vary as a result of local conditions. Since these conditions are unknown to the manufacturer, you must satisfy yourself as to local requirements. Do not exceed maximum rates or apply later than directed above.

WARNING: Do not use in combination with organic mercury fungicides or with all sprays.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Manufactured by AGWAY Inc.
AC179E

Syracuse, New York



USDA Reg. No. 8590-179

A.R. 21 1967