



Net Contents  
50 lbs.

DITHANE M-45 10 D

For Control of Certain Rose, Chrysanthemum and Potato Diseases

ACTIVE INGREDIENTS:

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate 8.0%

In which the ingredients are


Manganese . . . . . 1.6%

Zinc . . . . . 0.2%

Ethylene bisdithiocarbamate ion ( $C_4H_6N_2S_4$ ) . . . . . 6.2%

INERT INGREDIENTS . . . . . 92.0%

\*Rohm and Haas, U.S. Patents 2,504,404; 2,974,156 and Patent Pending



**CAUTION KEEP OUT OF REACH OF CHILDREN.**

May cause irritation of nose, throat and skin. Do not breathe dust. In case of skin contact, wash with water; for eyes, get medical attention.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse container. Destroy when empty.

IMPORTANT: Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of this product. Keep container closed when not in use.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and all not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

For best results apply when air is calm, covering foliage with a light film of dust. Apply as a preventive program making the first application before disease appears. Repeat applications as necessary. Keep new growth protected. This product is suitable for application through ground dusters or hand equipment.

**ROSES:** For control of Black Spot. Apply 10 to 15 lbs. per acre or  $\frac{1}{4}$  lb. per 1,000 square feet. Make the first application when the plants are well leafed and repeat at weekly intervals throughout the growing season.

**CHRYSANTHEMUMS:** For control of Botrytis Petal Spot. Apply at 10 to 20 lbs. per acre or  $\frac{1}{4}$  to  $\frac{1}{2}$  lbs. per 1,000 sq. feet before disease appears and repeat as necessary. Apply twice weekly during the blooming season.

**POTATOES:** For control of Early and Late Blight. Apply 10 to 20 lbs. per acre. Use lower rate when vines are small, increasing as vines become larger. Apply before disease appears and repeat at 7 to 14 day intervals throughout the growing season.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Agway, Inc. - Chemical Division, Box 1322, Syracuse, New York 13201

EPA Reg. No. 8590-182